



# 2018 Tentative/Scheduled Studies

## Residue and Efficacy/Crop Safety (E/CS)

(Order by Crop Group, Commodity, Chemical)

Print Date: 2/22/2018

<u>PR #</u>	<u>LAB</u>	<u>PRIORITY</u>	<u>STUDY DIRECTOR</u>	<u>CHEMICAL (MFG)</u>	<u>COMMODITY</u>	<u>CROP GROUP</u>
12351	18-MIR10	A	LENNON	BENZOVINDIFLUPYR + DIFENOCONAZOLE (SYNGEN)	BEET (GARDEN)	ROOT VEGETABLES SUBGROUPS (01AB)

**Reason for Need:** CERCOSPORA LEAF SPOT; LOSS OF LEAVES COMPROMISES ABILITY TO MACHINE HARVEST WHICH IS DONE BY PULLING LEAVES; DISEASE IS NOT BEING MANAGED ADEQUATELY WITH LABELED FUNGICIDES

**Use Pattern: (PCR):** USE THE APROVIA TOP DUAL AI PRODUCT; MAKE FOLIAR APPLIC OF 10.5-13.5 FL OZ/A; MAXIMUM 80 FL OZ/A/YEAR

**E/CS Data Requirements:**

**E/CS Research Comments:**

**IR-4 Residue Trial Plan:** JOINT PROJECT WITH PMC; NAFTA TRIAL SITES: 1 5-2 10 (ROOTS AND TOPS); 1 TRIAL IS A DECLINE

**Comments:** DIFENOCONAZOLE TOLERANCES EXIST FOR SUGARBEET, CARROT, GINSENG AND SUBGROUP 1C, AND MFG RESIDUE WORK IS ONGOING ON THE OTHER SUBGROUP 1B REP CROP RADISH; BENZOVINDIFLUPYR TOLERANCE EXISTS FOR SUBGROUP 1C:08/17; MFG SUPPORTS, RESIDUE ONLY, AND SUGGESTS DATA BE COLLECTED ON ROOTS AND TOPS FOR BOTH AI'S:09/17; POTENTIAL JOINT PROJECT WITH CANADA:10/17

**NER-EPA Region-FRD**

18-NJ246 Fisher, Jennifer

**NCR-EPA Region-FRD**

18-OH275 Doohan, D.

**SOR-EPA Region-FRD**

**WSR-EPA Region-FRD**

18-CA\*85 Benzen, Ms. Sharon D.  
(decline)  
18-CA86 Skiles, Keri

**CANADA-EPA Region-FRD**

18-ON287 Riddle, Geoff  
18-QC316 Cloutier, Dominic



# 2018 Tentative/Scheduled Studies

## Residue and Efficacy/Crop Safety (E/CS)

(Order by Crop Group, Commodity, Chemical)

Print Date: 2/22/2018

<u>PR #</u>	<u>LAB</u>	<u>PRIORITY</u>	<u>STUDY DIRECTOR</u>	<u>CHEMICAL (MFG)</u>	<u>COMMODITY</u>	<u>CROP GROUP</u>
12341	18-MIR08	A	LENNON	PROPICONAZOLE (ADAMA,SYNGEN)	GINSENG	ROOT VEGETABLES SUBGROUPS (01AB)

**Reason for Need:** FUSARIUM, ALTERNARIA - NEEDED DUE TO ALTERNARIA RESISTANCE TO BOSCALID AND STROBILURINS, AND NEED PRODUCTS ACTIVE AGAINST FUSARIUM

**Use Pattern: (PCR):** REQUESTED USE PATTERN IS SIMILAR TO LABELED USE ON CARROT AND RADISH (EXCEPT FOR RETREATMENT INTERVAL); USE THE TILT EC PRODUCT; MAKE 4-6 FOLIAR APPLIC OF 4 FL OZ/A, 14-DAY INTERVAL, 14-DAY PHI; USE 30-50 GPA FOR ALTERNARIA AND UP TO 100 GPA FOR FUSARIUM

**E/CS Data Requirements:**

**E/CS Research Comments:**

**IR-4 Residue Trial Plan:** 5-4; 1 TRIAL IS A DECLINE (1 TRIAL COVERED BY TASC \$\$)

**Comments:** THERE ARE TOLERANCES ON CARROT AND RADISH ROOTS; THE TILT LABEL IS THE SAME FOR CARROT AND RADISH; IR-4 CAN EXPLORE COVERING GINSENG TOLERANCE WITH A SUBGROUP 1B TOLERANCE, WITHOUT DOING RESIDUE WORK:08/17; MFG SUPPORTS, RESIDUE ONLY:09/17; MFG REQUESTS RESIDUE DATA BE GENERATED TO SUPPORT EXPORTS:01/18

**NER-EPA Region-FRD**

**NCR-EPA Region-FRD**

**SOR-EPA Region-FRD**

**WSR-EPA Region-FRD**

**CANADA-EPA Region-FRD**

- 18-WI204 Hausbeck, Dr. Mary K.  
(TASC \$)WI
- 18-WI205 Hausbeck, Dr. Mary K.  
WI
- 18-WI206 Hausbeck, Dr. Mary K.  
WI
- 18-WI207 Hausbeck, Dr. Mary K.  
WI (decline)



# 2018 Tentative/Scheduled Studies

## Residue and Efficacy/Crop Safety (E/CS)

(Order by Crop Group, Commodity, Chemical)

Print Date: 2/22/2018

<u>PR #</u>	<u>LAB</u>	<u>PRIORITY</u>	<u>STUDY DIRECTOR</u>	<u>CHEMICAL (MFG)</u>	<u>COMMODITY</u>	<u>CROP GROUP</u>
11912	18-TBD-CAN	A	CANADA	PYDIFLUMETOFEN (FTH 545) (SYNGEN)	GINSENG	ROOT VEGETABLES SUBGROUPS (01AB)

**Reason for Need:** ALTERNARIA, RHIZOCTONIA, FUSARIUM, CYLINDROCARPON RUSTY ROOT (IS A PRIORITY FOR THE INDUSTRY, AND THIS PRODUCT HAS SHOWN EFFICACY; MORE PRODUCTS ARE NEEDED TO COVER THE LONG GROWING PERIOD)

**Use Pattern: (PCR):** MAKE 4 APPLIC OF 75 GR AI/A, IN 30 GPA (FOLIAR) OR 100 GPA (SOIL APPLIC), 14-DAY INTERVAL, 30-DAY PHI; MFG IS DOING RESIDUE WORK WITH 75 G AI/HA, 4 APPLIC, 7-DAY INTERVAL, 7-DAY PHI, ON CROP SUBGROUP 1A:07/16; FROM 08/17 REQUEST: USE THE MIRAVIS PRODUCT; MAKE 4-6 FOLIAR APPLIC IN ENOUGH WATER TO PENETRATE INTO THE TOP SOIL LAYER, 14-DAY INTERVAL, 14-DAY PHI

**E/CS Data Requirements:** MFG REQUIRES COMPARATIVE PERFORMANCE DATA VS REGISTERED ALTERNATIVES WHERE LABELS ARE OF INTEREST:07/16

**E/CS Research Comments:**

**IR-4 Residue Trial Plan:** NEED 1 REG. 5 TRIAL (WI) IN 2018, ADDED TO CANADA PMC 2018 RESIDUE STUDY (NAFTA TRIAL SITES: 5-4)

**Comments:** KEY MARKETS INCLUDE TAIWAN, CHINA, HONG KONG; MANY DISEASE CONTROL PRODUCTS ARE NEEDED DUE TO THE LONG GROWING SEASON AND WEATHER CONDUCIVE TO DISEASE DEVELOPMENT; PERFORMANCE IS BEING TESTED IN MI IN 2016:12/15; MFG CONSIDERS THIS STILL UNDER EVALUATION, PENDING ADDITIONAL TECHNICAL OR OTHER INFORMATION ABOUT THE NEED FOR THIS USE FOR CONSIDERATION:05/16; MFG IS CONDUCTIING RESIDUE TRIALS ON REP CROPS RADISH, CARROT, SUGARBEET WHICH COULD COVER SUBGROUP 1A TOLERANCES, INCLUDING GINSENG; EFFICACY DATA WILL BE NEEDED FROM STATES WHERE LABELING IS NEEDED:07/16; IS AN "A" PRIORITY FOR RESIDUES IN 2018 IN CANADA, FOR A DRENCH USE PATTERN:05/17; HQ INCLUDING THIS AS A "RED A" TO ADD ONE U.S. TRIAL TO THE 2018 CANADIAN STUDY IN REGION 5 (WI) TO SUPPORT THE DRENCH USE PATTERN, AND TO SUPPORT THIS AS A JOINT NAFTA STUDY:09/17

**NER-EPA Region-FRD**

**NCR-EPA Region-FRD**

**SOR-EPA Region-FRD**

**WSR-EPA Region-FRD**

**CANADA-EPA Region-FRD**

18-MI190 Hausbeck, Dr. Mary K.

18-K\*169 TBD-CANADA  
18-K\*170 TBD-CANADA  
18-ON281 Wismer, R.J.



## 2018 Tentative/Scheduled Studies

### Residue and Efficacy/Crop Safety (E/CS)

*(Order by Crop Group, Commodity, Chemical)*

*Print Date: 2/22/2018*

<u>PR #</u>	<u>LAB</u>	<u>PRIORITY</u>	<u>STUDY DIRECTOR</u>	<u>CHEMICAL (MFG)</u>	<u>COMMODITY</u>	<u>CROP GROUP</u>
P11667	-		KUNKEL	BICYCLOPYRONE (SYNGEN)	HORSERADISH	ROOT VEGETABLES SUBGROUPS (01AB)

**Reason for Need:** SMALL-SEEDED BROADLEAF WEEDS, SUCH AS GLYPHOSATE RESISTANT PALMER AMARANTH AND COMMON WATERHEMP

**Use Pattern: (PCR):** MAKE ONE PRE-EMERGENCE SOIL APPLIC OF 2.56-3.42 FL OZ/A; PHI NOT DEFINED; NOT FOR FOLIAR USE

**E/CS Data Requirements:** MFG REQUESTS E/CS DATA FROM WI AND CA:09/19/16; MFG REQUESTS 5-6 TRIALS OVER 2 YEARS WITH 3 TRIALS IN YEAR 1, 2 STANDARD RATES (<50 G A

**E/CS Research Comments:**

**IR-4 Residue Trial Plan:**

**Comments:** FIRST USES REGISTERED BY EPA APRIL 2015, BOTH IN 4-WAY AI MIX AND SOLO PRODUCT:04/15; MFG OBJECTIVE FOR RESIDUES, BUT E/CS DATA NEEDED FROM WI AND CA:09/19/16

**NER-EPA Region-FRD**

**NCR-EPA Region-FRD**

**SOR-EPA Region-FRD**

**WSR-EPA Region-FRD**

**CANADA-EPA Region-FRD**

18-WIP01 Heider, Daniel J.  
18-WIP02 Heider, Daniel J.

18-CAP01 Hanson, Brad  
18-CAP02 Hanson, Brad



## 2018 Tentative/Scheduled Studies Residue and Efficacy/Crop Safety (E/CS)

*(Order by Crop Group, Commodity, Chemical)*

*Print Date: 2/22/2018*

<b>PR #</b>	<b>LAB</b>	<b>PRIORITY</b>	<b>STUDY DIRECTOR</b>	<b>CHEMICAL (MFG)</b>	<b>COMMODITY</b>	<b>CROP GROUP</b>
P8757	-	A	HOMA	FAMOXADONE + CYMOXANIL (DUPONT)	RADISH	ROOT VEGETABLES SUBGROUPS (01AB)

**Reason for Need:** FOLIAR DISEASE, ALTERNARIA, DOWNY MILDEW, WHITE RUST, CERCOSPORA

**Use Pattern: (PCR):** 1.5-2.5 LB PRODUCT/A; 30-50 GPA; 7-DAY RE-TREATMENT INTERVAL; 3-DAY PHI; 12-HOUR REI

**E/CS Data Requirements:** MFG CONFIRMED NEED FOR PERFORMANCE DATA BEFORE USE CAN BE LABELED:05/17; NEED AT LEAST 2 SUCCESSFUL TRIALS EACH FOR ALTERNARIA,

**E/CS Research Comments:**

**IR-4 Residue Trial Plan:** COMPLETE

**Comments:** CARROT IS DONE, BUT NOT SUGARBEET; CONSIDER AS 2018 "RED A" FOR E/CS DATA:05/17; THE RESIDUE STUDY SAMPLES WERE RECENTLY ANALYZED AND STORAGE STABILITY RESULTS WERE POOR FOR FAMOXADONE IN THE ROOTS; THE STUDY IS BEING REPEATED, UNDER PR# 12415; STUDY 08757 IS NOT BEING CANCELED:10/17; PERFORMANCE PROTOCOL SIGNED 2/12/18 AND E/CS DATA ON-GOING:02/18

**NER-EPA Region-FRD**

**NCR-EPA Region-FRD**

**SOR-EPA Region-FRD**

**WSR-EPA Region-FRD**

**CANADA-EPA Region-FRD**

18-MIP01 Hausbeck, Dr. Mary K.  
18-OHP01 Miller, Dr. Sally Ann

18-FLP01 Vallad, Gary

18-ORP01 Ocamb, Cindy



## 2018 Tentative/Scheduled Studies

### Residue and Efficacy/Crop Safety (E/CS)

*(Order by Crop Group, Commodity, Chemical)*

*Print Date: 2/22/2018*

<u>PR #</u>	<u>LAB</u>	<u>PRIORITY</u>	<u>STUDY DIRECTOR</u>	<u>CHEMICAL (MFG)</u>	<u>COMMODITY</u>	<u>CROP GROUP</u>
12415	18-CAR16	A	HOMA	FAMOXADONE + CYMOXANIL (DUPONT)	RADISH	ROOT VEGETABLES SUBGROUPS (01AB)

**Reason for Need:** FOLIAR DISEASE, ALTERNARIA, DOWNY MILDEW, WHITE RUST, CERCOSPORA

**Use Pattern: (PCR):** FROM PR# 08757: 1.5-2.5 LB PRODUCT/A; 30-50 GPA; 7-DAY RE-TREATMENT INTERVAL; 3-DAY PHI; 12-HOUR REI

**E/CS Data Requirements:** FROM PR# 08757: MFG CONFIRMED NEED FOR PERFORMANCE DATA BEFORE USE CAN BE LABELED:05/17

**E/CS Research Comments:**

**IR-4 Residue Trial Plan:** 1 3-2 5 10

**Comments:** THE PR# 08757 RESIDUE STUDY SAMPLES WERE ANALYZED AND STORAGE STABILITY RESULTS WERE POOR FOR FAMOXADONE IN THE ROOTS; THE STUDY NEEDS TO BE REPEATED IN 2018 UNDER THIS NEW PR#:10/17; RESEARCHABLE, RESIDUE & E/CS DATA NEEDED

**NER-EPA Region-FRD**

18-NJ251 Fisher, Jennifer

**NCR-EPA Region-FRD**

18-OH\*277 Horst, Leona

**SOR-EPA Region-FRD**

18-FL137 Thomas, Darrell  
(decline)  
18-GA\*147 Fraelich, Ben

**WSR-EPA Region-FRD**

18-CA102 Skiles, Keri

**CANADA-EPA Region-FRD**



# 2018 Tentative/Scheduled Studies

## Residue and Efficacy/Crop Safety (E/CS)

(Order by Crop Group, Commodity, Chemical)

Print Date: 2/22/2018

---

<u>PR #</u>	<u>LAB</u>	<u>PRIORITY</u>	<u>STUDY DIRECTOR</u>	<u>CHEMICAL (MFG)</u>	<u>COMMODITY</u>	<u>CROP GROUP</u>
P10558	-	+	ARSENOVIC	GLUFOSINATE (BAYER,UPI)	SWEET POTATO	TUBEROUS AND CORM VEGETABLES SUBGROUPS (01CD)

**Reason for Need:** ANNUAL BROADLEAF WEEDS

**Use Pattern: (PCR):** PLANT BURNDOWN; 29-43 OZ/A; 1 APPLIC; PER AR ME-TOO, FOR PRE-PLANT BURNDOWN AND AN ALTERNATIVE FOR BANDED APPLIC IN ROW MIDDLES

**E/CS Data Requirements:** MFG REQUIRES ONLY CROP SAFETY DATA FIRST:09/16

**E/CS Research Comments:**

**IR-4 Residue Trial Plan:**

**Comments:** ORIGINAL REQUEST WAS REC'D 3/24/2010; MFG WILL REVISIT AFTER RE-REG REVIEW IS COMPLETED BY EPA:05/16; MFG QUESTIONS IF THIS REQUEST IS SIMILAR TO GLUFOSINATE USE ON POTATO AS A DESICANT (HARVEST AID); A PRE-SEASON BURNDOWN WOULD POSSIBLY FIT IN THE RISK CUP; IF USE PATTERN NEEDED IS THE SAME AS POTATO, THEN NO RESIDUE DATA ARE NEEDED AND THE POTATO TOLERANCE WOULD COVER THIS REQUEST (SAME IS TRUE FOR TARO, 09568):08/16; EPA CAUTION; MFG SUPPORTS AS POTENTIAL, AND NEEDS ONLY CROP SAFETY DATA FIRST:09/16; E/CS BEFORE RESIDUE - E/CS ON-GOING: 01/17

NER-EPA Region-FRD

NCR-EPA Region-FRD

SOR-EPA Region-FRD

WSR-EPA Region-FRD

CANADA-EPA Region-FRD

18-NCP01 Jennings, K.

18-CAP03 Stoddard, Scott



# 2018 Tentative/Scheduled Studies

## Residue and Efficacy/Crop Safety (E/CS)

(Order by Crop Group, Commodity, Chemical)

Print Date: 2/22/2018

<u>PR #</u>	<u>LAB</u>	<u>PRIORITY</u>	<u>STUDY DIRECTOR</u>	<u>CHEMICAL (MFG)</u>	<u>COMMODITY</u>	<u>CROP GROUP</u>
12118	18-MIR04	A	LENNON	AZOXYSTROBIN + FLUDIOXONIL + DIFENOCONAZOLE (SYNGEN)	SWEET POTATO (POST HARVEST)	TUBEROUS AND CORM VEGETABLES SUBGROUPS (01CD)

**Reason for Need:** RHIZOPUS SOFT ROT (RHIZOPUS STOLONIFER), BLACK ROT (CERATOCYSTIS FIMBRIATA) - NO EFFECTIVE OPTIONS FOR POSTHARVEST DISEASE CONTROL

**Use Pattern: (PCR):** USE THE STADIUM PRODUCT; SPRAY 1.0 FL OZ/2000 LB OF ROOTS DURING PACKING

**E/CS Data Requirements:**

**E/CS Research Comments:**

**IR-4 Residue Trial Plan:** ANY 4 TRIALS (ALL 3 AI); PROCESSING (WET PEEL, "CHIPS") (ALL 4 TRIALS COVERED BY TASC \$\$)

**Comments:** KEY EXPORT MARKETS INCLUDE CANADA AND EUROPE; WAS TESTED IN PPWS FOR POSTHARVEST FUNGICIDES, PR# 11848; USE PATTERN REQUESTED HERE IS THE SAME AS WHAT IS LABELED FOR POSTHARVEST USE ON POTATO - CONSIDER IF A LABEL AMENDMENT CAN BE USED TO REGISTER THIS PRODUCT ON SWEET POTATO:01/17; MFG NEEDS TO SEE PERFORMANCE DATA BEFORE OK FOR RESIDUES (INCLUDING PROCESSING):05/17; MFG SUPPORTS, RESIDUE ONLY; RESIDUE DATA NEEDED FOR EU EXPORTS:09/17

**NER-EPA Region-FRD**

**NCR-EPA Region-FRD**

**SOR-EPA Region-FRD**

**WSR-EPA Region-FRD**

**CANADA-EPA Region-FRD**

18-NC217 Batts, Roger B. (TASC \$)(PROCESSING)	18-CA54 Skiles, Keri (TASC \$)
18-TX342 Marconi, Cristina (TASC \$)	18-CA55 Skiles, Keri (TASC \$)





## 2018 Tentative/Scheduled Studies Residue and Efficacy/Crop Safety (E/CS)

(Order by Crop Group, Commodity, Chemical)

Print Date: 2/22/2018

<b><u>PR #</u></b>	<b><u>LAB</u></b>	<b><u>PRIORITY</u></b>	<b><u>STUDY DIRECTOR</u></b>	<b><u>CHEMICAL (MFG)</u></b>	<b><u>COMMODITY</u></b>	<b><u>CROP GROUP</u></b>
P11620	-	A	ARSENOVIC	FOMESAFEN (SYNGEN)	ONION	ONION BULB AND GREEN SUBGROUPS (03-07AB)

**Reason for Need:** ANNUAL BROADLEAF WEEDS; ALSO IN DIRECT-SEEDED ONIONS

**Use Pattern: (PCR):** USE THE REFLEX PRODUCT (2.0 LB AI/GAL); MAKE 1-2 POSTEMERGENCE FOLIAR (AFTER THE FULL 1-LEAF STAGE OF ONION) APPLIC OF 0.125-0.25 LB AI/A, 15-DAY RE-TREATMENT INTERVAL; 40-DAY PHI; DO NOT EXCEED 0.25 LB AI/A/YEAR; DO NOT USE REFLEX ON THE SAME SOIL MORE THAN ONCE IN 2 YEARS; MFG RECOMMENDS 1 APPLIC OF 0.25 LB AI/A WITH A 30-DAY PHI:02/16

**E/CS Data Requirements:** MFG NEEDS TO SEE SUFFICIENT CROP SAFTEY DATA FROM WHICH TO DETERMINE IF REGISTRATION ON ONIONS IS ULTIMATELY SUPPORTABLE:02/15

**E/CS Research Comments:** IN 2016 PERFORMANCE TRIALS, EVALUATING 0.125 AND 0.25 LB AI/A RATES, WITH AND WITHOUT ADJUVANT, APPLIED ONCE FOLIAR BROADCAST TO 2-LF STAGE ONIONS, AND APPLIED TWICE - FIRST AT THE 2-LF STAGE FOLLOWED BY THE SAME TREATMENT 14 DAYS LATER, ALL IN >35 GPA; CROP INJURY, WEED CONTROL AND CROP YIELD DATA WILL BE COLLECTED

**IR-4 Residue Trial Plan:** BULB: 1 5 6 8 10-2 11 12; GREEN: 3 5 10-2 12 (1 DECLINE ON GR ONION)

**Comments:** MFG INDICATES CROP ROTATION AND GEOGRAPHIC USE RESTRICTIONS WILL LIMIT USE ON ONIONS:02/15; GR ONION (SEE PR# 11857) RESIDUE TRIALS TO BE INCLUDED IN THIS ONION PROTOCOL; CROP FOR THIS REQUEST CHANGED FROM DRY BULB TO "ONION" TO INCLUDE BULB AND GREEN ONION AND COVER THE WHOLE CROP GROUP; SEE 11857 FOR GR ONION PERFORMANCE TRIAL DATA:01/16

**NER-EPA Region-FRD**

18-NYP01    Hoepting, Christine

**NCR-EPA Region-FRD**

**SOR-EPA Region-FRD**

**WSR-EPA Region-FRD**

18-ORP02    Felix, J.

**CANADA-EPA Region-FRD**



# 2018 Tentative/Scheduled Studies

## Residue and Efficacy/Crop Safety (E/CS)

(Order by Crop Group, Commodity, Chemical)

Print Date: 2/22/2018

<b>PR #</b>	<b>LAB</b>	<b>PRIORITY</b>	<b>STUDY DIRECTOR</b>	<b>CHEMICAL (MFG)</b>	<b>COMMODITY</b>	<b>CROP GROUP</b>
12322	18-TBD	A	HOMA	NMG787 (TBD)	ONION	ONION BULB AND GREEN SUBGROUPS (03-07AB)

**Reason for Need:** NEMATODES - THERE IS A LACK OF REGISTERED PRODUCTS FOR NEMATODE MANAGEMENT, AND NEMATODES ARE IMPACTING PRODUCTION IN MAJOR GROWING AREAS

**Use Pattern: (PCR):** APPLY AS A DRENCH OR DRIP TO THE SOIL AT 500 G AI/HA; ALSO CONSIDER USE AS A SEED TREATMENT

**E/CS Data Requirements:**

**E/CS Research Comments:** MFG DOING A 2018 PERFORMANCE PROGRAM:11/17

**IR-4 Residue Trial Plan:** BULB: 1 6 8 10-2 11 GREEN: 2 3 5 10 (1 DECLINE ON EACH TYPE); FOR JOINT PROJECT WITH PMC, NAFTA TRIAL SITES = BULB: 5-2 6 8 10-2 11-2; GREEN: 5-2 10-2 12

**Comments:** USE ON GARLIC WOULD BE COVERED BY THESE RESIDUE DATA; NEED TO CONDUCT SUFFICIENT TRIALS TO SUPPORT A CODEX MRL; MFG CAN ASSIST BY ANALYZING RESIDUE SAMPLES:08/17; MFG INDICATES RESIDUE AND E/CS DATA ARE NEEDED:09/17; POTENTIAL JOINT PROJECT WITH CANADA:10/17

**NER-EPA Region-FRD**

**NCR-EPA Region-FRD**

**SOR-EPA Region-FRD**

**WSR-EPA Region-FRD**

**CANADA-EPA Region-FRD**

18-MI203 (bulb)	Zandstra, Dr. Bernard H.	18-TX347 (bulb)	Marconi, Cristina	18-CA78 (bulb)(decline)	Skiles, Keri	18-BC6 (green)(decline)	Clodius, Markus
18-OH273 (green)	Doohan, D.			18-CA79 (green)	Skiles, Keri	18-ON284 (bulb)	Weber-Henricks, Marj
				18-CA*80 (green)	Benzen, Ms. Sharon D.		
				18-CA81 (bulb)	Leach, Nathan	18-ON285 (green)	Riddle, Geoff
				18-ID160 (bulb)	Meeks, Mr. Will		
				18-NM259 (bulb)(reg 8)	Hamilton, Cary		
				18-OR301 (green)	Sturman, Peter		
				18-WA*360 (bulb)	Harvey, John		
				18-ID361 (bulb)	Meeks, Mr. Will		



# 2018 Tentative/Scheduled Studies

## Residue and Efficacy/Crop Safety (E/CS)

(Order by Crop Group, Commodity, Chemical)

Print Date: 2/22/2018

<b><u>PR #</u></b>	<b><u>LAB</u></b>	<b><u>PRIORITY</u></b>	<b><u>STUDY DIRECTOR</u></b>	<b><u>CHEMICAL (MFG)</u></b>	<b><u>COMMODITY</u></b>	<b><u>CROP GROUP</u></b>
12131	18-MIR05	A	BARNEY	PROMETRYN (ADAMA,SYNGEN)	LEEK	ONION, GREEN SUBGROUP (03-07B)

**Reason for Need:** WEEDS; NEED A POST EMERGENT OPTION FOR WEED CONTROL

**Use Pattern: (PCR):** USE THE CAPAROL 4L PRODUCT; MAKE 2 FOLIAR APPLIC OF 1.6 LB AI/A, 10-DAY INTERVAL, 50-DAY PHI; APPLY TO WELL ESTABLISHED TRANSPLANTS THAT HAVE GROWN ONE NEWLY ESTABLISHED LEAF, FOLLOWED BY A 2ND APPLIC IN 10 DAYS

**E/CS Data Requirements:** NEED 4-5 TRIALS, 1X AND 2X RATES (AND MIXTURE FOR 1X) VS STANDARD; CROP SAFETY % AT 7, 21, 42 DAT; EFFICACY AT 28 AND 42 DAT; YIELD; RUN TRI

**E/CS Research Comments:**

**IR-4 Residue Trial Plan:** JOINT PROJECT WITH PMC; NAFTA TRIAL SITES: 5-2 10-2 12 (THESE ARE TRIAL SITES FOR GR. ONION); 1 TRIAL IS A DECLINE

**Comments:** MRL IS ESTABLISHED IN KEY EXPORT MARKET IN CANADA:03/17; SYNGENTA SUPPORTS; NEED E/CS DATA WITH RESIDUE; AS NEEDED, GENERATE ENOUGH DATA TO SECURE CODEX MRL IF COMMODITY IS EXPORTED:07/17; EPA CAUTION:08/17; POTENTIAL JOINT PROJECT WITH CANADA:10/17

**NER-EPA Region-FRD**

**NCR-EPA Region-FRD**

**SOR-EPA Region-FRD**

**WSR-EPA Region-FRD**

**CANADA-EPA Region-FRD**

18-MI192      Zandstra, Dr. Bernard H.

18-CA56      Watkins, S.  
 18-CA\*57      Benzen, Ms. Sharon D.  
 18-OR296      Sturman, Peter  
 (decline)

18-ON282      Riddle, Geoff  
 18-QC314      Cloutier, Dominic



# 2018 Tentative/Scheduled Studies Residue and Efficacy/Crop Safety (E/CS)

(Order by Crop Group, Commodity, Chemical)

Print Date: 2/22/2018

<b><u>PR #</u></b>	<b><u>LAB</u></b>	<b><u>PRIORITY</u></b>	<b><u>STUDY DIRECTOR</u></b>	<b><u>CHEMICAL (MFG)</u></b>	<b><u>COMMODITY</u></b>	<b><u>CROP GROUP</u></b>
P12131	-	A	BARNEY	PROMETRYN (ADAMA,SYNGEN)	LEEK	ONION, GREEN SUBGROUP (03-07B)

**Reason for Need:** WEEDS; NEED A POST EMERGENT OPTION FOR WEED CONTROL

**Use Pattern: (PCR):** USE THE CAPAROL 4L PRODUCT; MAKE 2 FOLIAR APPLIC OF 1.6 LB AI/A, 10-DAY INTERVAL, 50-DAY PHI; APPLY TO WELL ESTABLISHED TRANSPLANTS THAT HAVE GROWN ONE NEWLY ESTABLISHED LEAF, FOLLOWED BY A 2ND APPLIC IN 10 DAYS

**E/CS Data Requirements:** NEED 4-5 TRIALS, 1X AND 2X RATES (AND MIXTURE FOR 1X) VS STANDARD; CROP SAFETY % AT 7, 21, 42 DAT; EFFICACY AT 28 AND 42 DAT; YIELD; RUN TRI

**E/CS Research Comments:**

**IR-4 Residue Trial Plan:** JOINT PROJECT WITH PMC; NAFTA TRIAL SITES: 5-2 10-2 12 (THESE ARE TRIAL SITES FOR GR. ONION); 1 TRIAL IS A DECLINE

**Comments:** MRL IS ESTABLISHED IN KEY EXPORT MARKET IN CANADA:03/17; SYNGENTA SUPPORTS; NEED E/CS DATA WITH RESIDUE; AS NEEDED, GENERATE ENOUGH DATA TO SECURE CODEX MRL IF COMMODITY IS EXPORTED:07/17; EPA CAUTION:08/17; POTENTIAL JOINT PROJECT WITH CANADA:10/17

**NER-EPA Region-FRD**

**NCR-EPA Region-FRD**

**SOR-EPA Region-FRD**

**WSR-EPA Region-FRD**

**CANADA-EPA Region-FRD**

18-MIP02	Zandstra, Dr. Bernard H.	18-FLP02	Dittmar, Dr. Peter	18-CAP04	Smith, Dr. Richard
----------	--------------------------	----------	--------------------	----------	--------------------



## 2018 Tentative/Scheduled Studies

### Residue and Efficacy/Crop Safety (E/CS)

*(Order by Crop Group, Commodity, Chemical)*

*Print Date: 2/22/2018*

<b><u>PR #</u></b>	<b><u>LAB</u></b>	<b><u>PRIORITY</u></b>	<b><u>STUDY DIRECTOR</u></b>	<b><u>CHEMICAL (MFG)</u></b>	<b><u>COMMODITY</u></b>	<b><u>CROP GROUP</u></b>
P11857	-		ARSENOVIC	FOMESAFEN (SYNGEN)	ONION (GREEN)	ONION, GREEN SUBGROUP (03-07B)

**Reason for Need:** ANNUAL BROADLEAF WEEDS (IN DIRECT-SEEDED GR ONIONS)

**Use Pattern: (PCR):** USE THE REFLEX PRODUCT; MAKE 1-2 POSTEMERGENCE FOLIAR (AFTER THE FULL 1-LEAF STAGE OF ONION) APPLIC OF 0.125-0.25 LB AI/A, 15-DAY INTERVAL, 30-DAY PHI; DO NOT EXCEED 0.25 LB AI/A/YEAR

**E/CS Data Requirements:**

**E/CS Research Comments:** IN 2016 PERFORMANCE TRIALS, EVALUATING 0.125/0.25/0.5 LB AI/A, OR 0.195/0.375/0.75 LB AI/A RATES AT PRE (WITHOUT ADJUVANT, AFTER SEEDING BUT BEFORE ONION EMERGENCE) AND FOLIAR BROADCAST (WITH ADJUVANT, 28 DAYS AFTER PLANTING) TIMINGS, ALL IN 10-20 GPA; CROP INJURY, WEED CONTROL AND CROP YIELD DATA WILL BE COLLECTED

**IR-4 Residue Trial Plan:** GREEN: 3 5 10-2 12

**Comments:** DEVELOPMENT OF RESIDUE DATA CAN BE INCLUDED WITHIN BULB ONION PROJECT #11620:12/15; MFG MADE GREEN ONION RESEARCHABLE; RESIDUE TRIALS INCLUDED WITH BULB ONION TRIALS IN PROTOCOL 11620, BUT PERFORMANCE TRIALS ON GR ONION ARE CAPTURED WITH THIS GR ONION PR:01/16

**NER-EPA Region-FRD**

**NCR-EPA Region-FRD**

**SOR-EPA Region-FRD**

**WSR-EPA Region-FRD**

**CANADA-EPA Region-FRD**

18-SC\*P01 Wade, Paul

18-ORP03 Peachey, Ed



## 2018 Tentative/Scheduled Studies

### Residue and Efficacy/Crop Safety (E/CS)

*(Order by Crop Group, Commodity, Chemical)*

*Print Date: 2/22/2018*

<b>PR #</b>	<b>LAB</b>	<b>PRIORITY</b>	<b>STUDY DIRECTOR</b>	<b>CHEMICAL (MFG)</b>	<b>COMMODITY</b>	<b>CROP GROUP</b>
12292	18-FLR06	A	LENNON	SPINETORAM+SULFOXAFLO R (DOWAGR)	LETTUCE (GH)	LEAFY GREENS SUBGROUP (04-16A)

**Reason for Need:** APHIDS, WHITEFLIES, LEPIDOPTERA

**Use Pattern: (PCR):** USE THE XXPIRE WG PRODUCT, AND USE RADIANT AI/A RATES; MAKE 3-4 FOLIAR APPLIC, 10-14 DAY INTERVALS, 0-1 DAY PHI

**E/CS Data Requirements:**

**E/CS Research Comments:**

**IR-4 Residue Trial Plan:** ANY 4 TRIALS (2 HEAD, 2 LEAF); 1 LEAF TRIAL IS A DECLINE; FOR JOINT PROJECT WITH PMC, NAFTA TRIAL SITES: ANY 4

**Comments:** CANADA IS A KEY EXPORT MARKET; CAN COVER SOLO SPINETORAM REQUEST PR#12037 WITH THIS DUAL AI PROJECT:07/17; MFG SUPPORTS, RESIDUE ONLY, AND DESIRES THAT RESIDUES SUPPORT CURRENT TOLERANCES SO THERE'S NO NEED TO REQUEST ANY INCREASE:09/17; POTENTIAL JOINT PROJECT WITH CANADA:10/17

**NER-EPA Region-FRD**

18-MD177    Ross, Marylee  
(leaf)

**NCR-EPA Region-FRD**

18-OH\*271    Horst, Leona  
(leaf)

**SOR-EPA Region-FRD**

18-TX346    Marconi, Cristina  
(leaf)(decline)

**WSR-EPA Region-FRD**

18-CA75    Ennes, D. (Kearney)  
(head)  
18-CA76    Ennes, D. (Kearney)  
(leaf)

**CANADA-EPA Region-FRD**

18-BC5    Nield, David  
(head)



# 2018 Tentative/Scheduled Studies Residue and Efficacy/Crop Safety (E/CS)

(Order by Crop Group, Commodity, Chemical)

Print Date: 2/22/2018

<u>PR #</u>	<u>LAB</u>	<u>PRIORITY</u>	<u>STUDY DIRECTOR</u>	<u>CHEMICAL (MFG)</u>	<u>COMMODITY</u>	<u>CROP GROUP</u>
12388	18-CAR15	A	HOMA	FLUTIANIL (LANDIS,OATAGRIO)	LETTUCE (HEAD & LEAF)	LEAFY GREENS SUBGROUP (04-16A)

**Reason for Need:** POWDERY MILDEW

**Use Pattern: (PCR):** FROM 09180 REQUEST: 0.04 LB AI/A; MAX OF 3 APPLIC/SEASON; FROM 09180 RESIDUE PROTOCOL: 0.04 LB AI/A, MAKE 5 FOLIAR APPLIC IN 30-80 GPA AT 7-DAY INTERVALS, 0-DAY PHI

**E/CS Data Requirements:** MFG REQUIRES SOME GH PERFORMANCE DATA:09/17

**E/CS Research Comments:**

**IR-4 Residue Trial Plan:** HEAD: 1/2 3 10-4 LEAF: 1/2 3 10-4 GH: ANY 4 TRIALS (2 HEAD, 2 LEAF); 1 FIELD/LEAF TRIAL AND 1 GH/LEAF TRIAL ARE DECLINES

**Comments:** ORIG PCR (09180) RECD 03/13/04; MFG WILL NOT SUPPORT:08/09; HOLD AS THERE ARE NO FOOD TOLERANCES YET:06/15; MFG ASSESSMENT OF 2007 STUDY DATA (09180) INDICATES IT CAN NOT BE SUBMITTED; MFG RE-EVALUATING THE REQUEST:08/16; MFG CONFIRMED THIS REQUEST IS NOW RESEARCHABLE:08/25/16; SINCE PREFIX "A" PR#'S ARE NO LONGER ASSIGNED, DATA TO THIS POINT WAS COPIED FROM RELATED 2007 STUDY, PR# 09180; MFG ALSO SUPPORTS INCLUSION OF GH USE (PR# 12374), BUT WITH SOME PERFORMANCE DATA:09/17

[NER-EPA Region-FRD](#)

[NCR-EPA Region-FRD](#)

[SOR-EPA Region-FRD](#)

[WSR-EPA Region-FRD](#)

[CANADA-EPA Region-FRD](#)



## 2018 Tentative/Scheduled Studies

### Residue and Efficacy/Crop Safety (E/CS)

*(Order by Crop Group, Commodity, Chemical)*

*Print Date: 2/22/2018*

18-MD180 (gh leaf)	Ross, Marylee	18-WI387 (gh leaf)	Chapman, Scott	18-FL133 (gh head)	Long, Michael	18-CA87 (field leaf)	Kyser, Guy
18-NJ247 (field head)	Fisher, Jennifer			18-FL134 (field head)(reg. 3)	Thomas, Darrell	18-CA88 (field head)	Kyser, Guy
18-NJ248 (field leaf)	Fisher, Jennifer			18-FL135 (field leaf)(reg. 3)	Long, Michael	18-CA89 (field leaf)	Leach, Nathan
				18-SC*329 (field leaf)	Wade, Paul	18-CA90 (field head)	Leach, Nathan
				18-TX348 (gh head)	Marconi, Cristina	18-CA91 (field leaf)	Skiles, Keri
						18-CA92 (gh leaf)(decline)	Ennes, D. (Kearney)
						18-CA93 (field head)	Skiles, Keri
						18-CA*94 (field leaf)(decline)	Benzen, Ms. Sharon D.
						18-CA*95 (field head)	Benzen, Ms. Sharon D.
						18-NM260 (field head)	Hamilton, Cary
						18-NM261 (field leaf)	Hamilton, Cary





# 2018 Tentative/Scheduled Studies Residue and Efficacy/Crop Safety (E/CS)

(Order by Crop Group, Commodity, Chemical)

Print Date: 2/22/2018

<b><u>PR #</u></b>	<b><u>LAB</u></b>	<b><u>PRIORITY</u></b>	<b><u>STUDY DIRECTOR</u></b>	<b><u>CHEMICAL (MFG)</u></b>	<b><u>COMMODITY</u></b>	<b><u>CROP GROUP</u></b>
P12388	-	A	HOMA	FLUTIANIL (LANDIS,OATAGRIO)	LETTUCE (HEAD & LEAF)	LEAFY GREENS SUBGROUP (04-16A)

**Reason for Need:** POWDERY MILDEW

**Use Pattern: (PCR):** FROM 09180 REQUEST: 0.04 LB AI/A; MAX OF 3 APPLIC/SEASON; FROM 09180 RESIDUE PROTOCOL: 0.04 LB AI/A, MAKE 5 FOLIAR APPLIC IN 30-80 GPA AT 7-DAY INTERVALS, 0-DAY PHI

**E/CS Data Requirements:** MFG REQUIRES SOME GH PERFORMANCE DATA:09/17

**E/CS Research Comments:**

**IR-4 Residue Trial Plan:** HEAD: 1/2 3 10-4 LEAF: 1/2 3 10-4 GH: ANY 4 TRIALS (2 HEAD, 2 LEAF); 1 FIELD/LEAF TRIAL AND 1 GH/LEAF TRIAL ARE DECLINES

**Comments:** ORIG PCR (09180) RECD 03/13/04; MFG WILL NOT SUPPORT:08/09; HOLD AS THERE ARE NO FOOD TOLERANCES YET:06/15; MFG ASSESSMENT OF 2007 STUDY DATA (09180) INDICATES IT CAN NOT BE SUBMITTED; MFG RE-EVALUATING THE REQUEST:08/16; MFG CONFIRMED THIS REQUEST IS NOW RESEARCHABLE:08/25/16; SINCE PREFIX "A" PR#'S ARE NO LONGER ASSIGNED, DATA TO THIS POINT WAS COPIED FROM RELATED 2007 STUDY, PR# 09180; MFG ALSO SUPPORTS INCLUSION OF GH USE (PR# 12374), BUT WITH SOME PERFORMANCE DATA:09/17

**NER-EPA Region-FRD**

18-NYP02     McGrath, Dr. Margaret  
(GH)

**NCR-EPA Region-FRD**

**SOR-EPA Region-FRD**

18-FLP03     Vallad, Gary  
(GH)

**WSR-EPA Region-FRD**

18-AZP01     Matheron, Dr. Michael E.  
(GH)

**CANADA-EPA Region-FRD**



# 2018 Tentative/Scheduled Studies

## Residue and Efficacy/Crop Safety (E/CS)

(Order by Crop Group, Commodity, Chemical)

Print Date: 2/22/2018

---

<u>PR #</u>	<u>LAB</u>	<u>PRIORITY</u>	<u>STUDY DIRECTOR</u>	<u>CHEMICAL (MFG)</u>	<u>COMMODITY</u>	<u>CROP GROUP</u>
11027	17-CAR06	A	JOLLY	MANDESTROBIN (S-2200) (VALENT)	LETTUCE (HEAD & LEAF)	LEAFY GREENS SUBGROUP (04-16A)

**Reason for Need:** SCLEROTINIA MINOR, SCLEROTINIA SCLEROTIORUM (LETTUCE DROP)

**Use Pattern: (PCR):** 0.375 LB AI/A; 2 APPLIC, TO SOIL AND/OR FOLIAGE OF VERY YOUNG PLANTS

**E/CS Data Requirements:** CANADA CONDUCTED 3 EFFICACY TRIALS IN 2017

**E/CS Research Comments:**

**IR-4 Residue Trial Plan:** HEAD: 5-2 10-6, 1 DECLINE TRIAL; LEAF: 5-2 10-6, 1 DECLINE TRIAL (HEAD AND LEAF TRIALS = NAFTA SITES); NEED 1 "RED A" TRIAL IN 2018

**Comments:** MFG HOLD:08/12; FIRST REGISTRATION PENDING ON CORN (FIELD/POP/SWEET), LEGUME VEG SUCULENT OR DRY, SUBGROUPS 13-07F & 13-07G, RAPESEED, TURF; CHANGED STATUS TO UNDER EVAL:03/16; MFG SUPPORTS, NO E/CS DATA NEEDED:05/16; REQUESTOR ASKED FOR THIS TO COVER LETTUCE (COMMODITY CHANGED TO LETTUCE/HEAD & LEAF); CANADA INTEREST:09/16; IS A JOINT PROJECT WITH CANADA:05/17

**NER-EPA Region-FRD**

**NCR-EPA Region-FRD**

**SOR-EPA Region-FRD**

**WSR-EPA Region-FRD**

**CANADA-EPA Region-FRD**

18-MI187      Zandstra, Dr. Bernard H.  
(head)

18-NM392      Hamilton, Cary  
(head)



## 2018 Tentative/Scheduled Studies

### Residue and Efficacy/Crop Safety (E/CS)

*(Order by Crop Group, Commodity, Chemical)*

*Print Date: 2/22/2018*

<u>PR #</u>	<u>LAB</u>	<u>PRIORITY</u>	<u>STUDY DIRECTOR</u>	<u>CHEMICAL (MFG)</u>	<u>COMMODITY</u>	<u>CROP GROUP</u>
12029	17-FLR06	A	BARNEY	PROMETRYN (ADAMA,SYNGEN)	SPINACH	LEAFY GREENS SUBGROUP (04-16A)

**Reason for Need:** REDUCE THE PLANT BACK RESTRICTION FROM 12 MONTHS TO 3-4 MONTHS FOR SPINACH, FOLLOWING USE ON CILANTRO PER REGISTERED CAPAROL LABEL

**Use Pattern: (PCR):** USE CAPAROL 4L PRODUCT PER REGISTERED LABEL USE DIRECTIONS FOR WEED CONTROL IN CILANTRO

**E/CS Data Requirements:** MFG REQUIRES MINIMUM OF 2 CROP SAFETY TRIALS WITH SPINACH, PREFERABLY IN CA, WITH EVALUATIONS AT THE DESIRED 3-MONTH PLANTBACK IN

**E/CS Research Comments:**

**IR-4 Residue Trial Plan:** 1 2-2 6-2 9 10-2; NEED 2 "RED A" TRIALS IN 2018

**Comments:** SEE ALSO PROJECT REQUESTS 12034/CABBAGE (FOR REQUESTED NAPA CABAGE), 12035/BROCCOLI (FOR REQUESTED BRUSSELS SPROUTS) AND 12036/BELL AND NON-BELL PEPPER WHICH WERE ALL PART OF THE ORIGINAL REQUEST:08/16; MFG SUPPORTS FOR RESIDUE WORK, BUT SIMULTANEOUS SPINACH PLANTBACK CROP SAFETY TRIALS ARE REQUIRED:09/16

**NER-EPA Region-FRD**

**NCR-EPA Region-FRD**

**SOR-EPA Region-FRD**

**WSR-EPA Region-FRD**

**CANADA-EPA Region-FRD**

18-SC\*323 Wade, Paul  
18-SC\*324 Wade, Paul



## 2018 Tentative/Scheduled Studies

### Residue and Efficacy/Crop Safety (E/CS)

*(Order by Crop Group, Commodity, Chemical)*

*Print Date: 2/22/2018*

<b><u>PR #</u></b>	<b><u>LAB</u></b>	<b><u>PRIORITY</u></b>	<b><u>STUDY DIRECTOR</u></b>	<b><u>CHEMICAL (MFG)</u></b>	<b><u>COMMODITY</u></b>	<b><u>CROP GROUP</u></b>
P12029	-	A	BARNEY	PROMETRYN (ADAMA,SYNGEN)	SPINACH	LEAFY GREENS SUBGROUP (04-16A)

**Reason for Need:** REDUCE THE PLANT BACK RESTRICTION FROM 12 MONTHS TO 3-4 MONTHS FOR SPINACH, FOLLOWING USE ON CILANTRO PER REGISTERED CAPAROL LABEL

**Use Pattern: (PCR):** USE CAPAROL 4L PRODUCT PER REGISTERED LABEL USE DIRECTIONS FOR WEED CONTROL IN CILANTRO

**E/CS Data Requirements:** MFG REQUIRES MINIMUM OF 2 CROP SAFETY TRIALS WITH SPINACH, PREFERABLY IN CA, WITH EVALUATIONS AT THE DESIRED 3-MONTH PLANTBACK IN

**E/CS Research Comments:**

**IR-4 Residue Trial Plan:** 1 2-2 6-2 9 10-2; NEED 2 "RED A" TRIALS IN 2018

**Comments:** SEE ALSO PROJECT REQUESTS 12034/CABBAGE (FOR REQUESTED NAPA CABAGE), 12035/BROCCOLI (FOR REQUESTED BRUSSELS SPROUTS) AND 12036/BELL AND NON-BELL PEPPER WHICH WERE ALL PART OF THE ORIGINAL REQUEST:08/16; MFG SUPPORTS FOR RESIDUE WORK, BUT SIMULTANEOUS SPINACH PLANTBACK CROP SAFETY TRIALS ARE REQUIRED:09/16

**NER-EPA Region-FRD**

**NCR-EPA Region-FRD**

**SOR-EPA Region-FRD**

**WSR-EPA Region-FRD**

**CANADA-EPA Region-FRD**

18-FLP04 Dittmar, Dr. Peter	18-CAP06 Fennimore, S.
(multi-crop: spinach, pepper, brussel sprout)	(spinach only)



## 2018 Tentative/Scheduled Studies

### Residue and Efficacy/Crop Safety (E/CS)

*(Order by Crop Group, Commodity, Chemical)*

*Print Date: 2/22/2018*

<b><u>PR #</u></b>	<b><u>LAB</u></b>	<b><u>PRIORITY</u></b>	<b><u>STUDY DIRECTOR</u></b>	<b><u>CHEMICAL (MFG)</u></b>	<b><u>COMMODITY</u></b>	<b><u>CROP GROUP</u></b>
P11519	-	+	ARSENOVIC	CLOMAZONE (FMC)	GREENS (MUSTARD)	BRASSICA LEAFY GREENS SUBGROUP (04-16B)

**Reason for Need:** COMMON PURSLANE, VELVETLEAF, PROSTRATE PIGWEED, ANNUAL GRASS; THERE ARE FEW OPTIONS FOR BRASSICA GREENS

**Use Pattern: (PCR):** 0.25-1.0 LB/A; ONE SOIL PREEMERGENCE APPLIC OF COMMAND 3 ME, AFTER SEEDING BUT BEFORE CROP EMERGENCE

**E/CS Data Requirements:** PER MFG, NEED ONE MORE CROP SAFETY TRIAL IN GA:09/17; MFG REVISION - PREFER AT LEAST 3 TRIALS OVER 2 YRS; 6 TRTS, 1 APPLIC:10/17

**E/CS Research Comments:**

**IR-4 Residue Trial Plan:**

**Comments:** EPA CAUTION:08/14; MFG NEEDS E/CS DATA BEFORE SUPPORTING RESIDUE STUDY:07/15

**NER-EPA Region-FRD**

**NCR-EPA Region-FRD**

**SOR-EPA Region-FRD**

**WSR-EPA Region-FRD**

**CANADA-EPA Region-FRD**

18-OHP02     Doohan, D.

18-GA\*P03     Fraelich, Ben

18-CAP07     Hanson, Brad  
18-WA\*P01     Harvey, John



# 2018 Tentative/Scheduled Studies Residue and Efficacy/Crop Safety (E/CS)

(Order by Crop Group, Commodity, Chemical)

Print Date: 2/22/2018

<u>PR #</u>	<u>LAB</u>	<u>PRIORITY</u>	<u>STUDY DIRECTOR</u>	<u>CHEMICAL (MFG)</u>	<u>COMMODITY</u>	<u>CROP GROUP</u>
07091	18-YAR01	A	BARNEY	FLUAZINAM (ISK,SYNGEN)	BROCCOLI	BRASSICA HEAD AND STEM VEGETABLE GROUP (05-16)

**Reason for Need:** FOLIAR DISEASES, DOWNY MILDEW, WIRESTEM (THE ONLY WIRESTEM FUNGICIDE CAN ONLY BE USED AT PLANTING)

**Use Pattern: (PCR):** FOLIAR SPRAY; 0.5 LB AI/A (SEE CABBAGE STUDIES 08796 AND 07093 FOR DRENCH AND FOLIAR USE PATTERNS, INCLUDING 7-DAY PHI)

**E/CS Data Requirements:** KEY IS DOWNY MILDEW; NEED 3 TRIALS, FOLIAR AND SOIL TRTS (?):10/17

**E/CS Research Comments:**

**IR-4 Residue Trial Plan:** 6 10-4 12 (1 DECLINE) -- THIS COMPLETES THE CROP GROUP

**Comments:** SYNGENTA & ISK SUPPORT:10/02; EARLY SEASON USE; THERE IS A CROP GROUP 5 TOLERANCE, BUT NOT COVERING FOLIAR USES:06/14; CLUB ROOT IS ON THE LABEL AND A SINGLE TRIAL FOR RHIZOCTONIA IS DONE:09/16

**NER-EPA Region-FRD**

**NCR-EPA Region-FRD**

**SOR-EPA Region-FRD**

**WSR-EPA Region-FRD**

**CANADA-EPA Region-FRD**

18-TX335 Marconi, Cristina

18-CA8 Leach, Nathan  
 18-CA9 Skiles, Keri  
 18-CA\*10 Benzen, Ms. Sharon D.  
 18-NM252 Hamilton, Cary  
 (reg 10)  
 18-OR288 Koskela, Ms. Gina  
 (reg 12)(decline)



## 2018 Tentative/Scheduled Studies

### Residue and Efficacy/Crop Safety (E/CS)

*(Order by Crop Group, Commodity, Chemical)*

*Print Date: 2/22/2018*

<b><u>PR #</u></b>	<b><u>LAB</u></b>	<b><u>PRIORITY</u></b>	<b><u>STUDY DIRECTOR</u></b>	<b><u>CHEMICAL (MFG)</u></b>	<b><u>COMMODITY</u></b>	<b><u>CROP GROUP</u></b>
P7091	-NONE	A	HOMA	FLUAZINAM (ISK,SYNGEN)	BROCCOLI	BRASSICA HEAD AND STEM VEGETABLE GROUP (05-16)

**Reason for Need:** FOLIAR DISEASES, DOWNY MILDEW, WIRESTEM (THE ONLY WIRESTEM FUNGICIDE CAN ONLY BE USED AT PLANTING)

**Use Pattern: (PCR):** FOLIAR SPRAY; 0.5 LB AI/A (SEE CABBAGE STUDIES 08796 AND 07093 FOR DRENCH AND FOLIAR USE PATTERNS, INCLUDING 7-DAY PHI)

**E/CS Data Requirements:** KEY IS DOWNY MILDEW; NEED 3 TRIALS, FOLIAR AND SOIL TRTS (?):10/17

**E/CS Research Comments:**

**IR-4 Residue Trial Plan:** 6 10-4 12 (1 DECLINE) -- THIS COMPLETES THE CROP GROUP

**Comments:** SYNGENTA & ISK SUPPORT:10/02; EARLY SEASON USE; THERE IS A CROP GROUP 5 TOLERANCE, BUT NOT COVERING FOLIAR USES:06/14; CLUB ROOT IS ON THE LABEL AND A SINGLE TRIAL FOR RHIZOCTONIA IS DONE:09/16

**NER-EPA Region-FRD**

**NCR-EPA Region-FRD**

**SOR-EPA Region-FRD**

**WSR-EPA Region-FRD**

**CANADA-EPA Region-FRD**

18-OHP03     Miller, Dr. Sally Ann

18-FLP05     Vallad, Gary

18-ORP04     Ocamb, Cindy



## 2018 Tentative/Scheduled Studies

### Residue and Efficacy/Crop Safety (E/CS)

*(Order by Crop Group, Commodity, Chemical)*

*Print Date: 2/22/2018*

<b>PR #</b>	<b>LAB</b>	<b>PRIORITY</b>	<b>STUDY DIRECTOR</b>	<b>CHEMICAL (MFG)</b>	<b>COMMODITY</b>	<b>CROP GROUP</b>
10557	18-FLR02	A	ARSENOVIC	SULFENTRAZONE (FMC)	BROCCOLI	BRASSICA HEAD AND STEM VEGETABLE GROUP (05-16)

**Reason for Need:** HOPHORNBEAN COPPERLEAF, PIGWEED

**Use Pattern: (PCR):** POST-TRANSPLANT DIRECTED BETWEEN THE ROWS; 2.25-12.0 OZ A/A; 1 SOIL APPLIC; 30-DAY PHI; PER ME-TOO REQUEST, NEED 30-DAY PHI, WHETHER FOR TRANSPLANTED OR DIRECT-SEEDED CROP:09/17

**E/CS Data Requirements:** MFG REQUESTS AN E/CS TRIAL IN GA, ON SEEDED CROP WITH APPLIC 14 DAYS AFTER PLANTING:06/17

**E/CS Research Comments:** IN 2017 PERFORMANCE PROTOCOL: USING SPARTAN 4F, IN NC AND FL, TEST 0.094, 0.1875, 0.375 AND 0.75 LB A/A VS A GROWER STANDARD, MAKING A SINGLE SHIELDED/HOODED TO ROW MIDDLES APPLIC 30 DAYS BEFORE FIRST HARVEST, IN AT LEAST 15 GPA, IN CALIBRATED EQUIPMENT THAT PROVIDES A UNIFORM SPRAY; EVALUATE CROP INJURY, WEED CONTROL AND YIELD; IN 2018 PROTOCOL: IN A SINGLE GA TRIAL, TEST THE SAME 4 RATE S OF SPARTAN 4F IN SAME GPA, BUT MAKE THE SHIELDED/HOODED APPLIC 14 DAYS AFTER SEEDING, AND MAKE THE SAME EVALUATIONS

**IR-4 Residue Trial Plan:** JOINT PROJECT WITH CANADA PMC; NAFTA TRIAL SITES: 5-2 10-6

**Comments:** MFG REQUIRES E/CS DATA PRIOR TO RESIDUE STUDY:09/12; TOLERANCE IS ESTABLISHED, PRE-PLANT USE IS REGISTERED:09/14; E/CS RESEARCH IS ONGOING, BUT NEED MORE POST-TRANSPLANT E/CS DATA BEFORE RESIDUE STUDY:06/17; MFG GAVE OK FOR RESIDUE STUDY AT FUW, SIMILAR TO CABBAGE (10556):09/17; POTENTIAL JOINT PROJECT WITH CANADA:10/17

**NER-EPA Region-FRD**

**NCR-EPA Region-FRD**

**SOR-EPA Region-FRD**

**WSR-EPA Region-FRD**

**CANADA-EPA Region-FRD**

18-CA30	Kyser, Guy	18-ON279	Weber-Henricks, Mary
18-CA31	Leach, Nathan	18-QC313	Cloutier, Dominic
18-CA32	Skiles, Keri		
18-CA*33	Benzen, Ms. Sharon D.		
18-CA*34	Benzen, Ms. Sharon D.		
18-NM254	Hamilton, Cary		





## 2018 Tentative/Scheduled Studies Residue and Efficacy/Crop Safety (E/CS)

(Order by Crop Group, Commodity, Chemical)

Print Date: 2/22/2018

<b><u>PR #</u></b>	<b><u>LAB</u></b>	<b><u>PRIORITY</u></b>	<b><u>STUDY DIRECTOR</u></b>	<b><u>CHEMICAL (MFG)</u></b>	<b><u>COMMODITY</u></b>	<b><u>CROP GROUP</u></b>
P10557	-NONE	A	ARSENOVIC	SULFENTRAZONE (FMC)	BROCCOLI	BRASSICA HEAD AND STEM VEGETABLE GROUP (05-16)

**Reason for Need:** HOPHORNBEAN COPPERLEAF, PIGWEED

**Use Pattern: (PCR):** POST-TRANSPLANT DIRECTED BETWEEN THE ROWS; 2.25-12.0 OZ A/A; 1 SOIL APPLIC; 30-DAY PHI; PER ME-TOO REQUEST, NEED 30-DAY PHI, WHETHER FOR TRANSPLANTED OR DIRECT-SEEDED CROP:09/17

**E/CS Data Requirements:** MFG REQUESTS AN E/CS TRIAL IN GA, ON SEEDED CROP WITH APPLIC 14 DAYS AFTER PLANTING:06/17

**E/CS Research Comments:** IN 2017 PERFORMANCE PROTOCOL: USING SPARTAN 4F, IN NC AND FL, TEST 0.094, 0.1875, 0.375 AND 0.75 LB A/A VS A GROWER STANDARD, MAKING A SINGLE SHIELDED/HOODED TO ROW MIDDLES APPLIC 30 DAYS BEFORE FIRST HARVEST, IN AT LEAST 15 GPA, IN CALIBRATED EQUIPMENT THAT PROVIDES A UNIFORM SPRAY; EVALUATE CROP INJURY, WEED CONTROL AND YIELD; IN 2018 PROTOCOL: IN A SINGLE GA TRIAL, TEST THE SAME 4 RATE S OF SPARTAN 4F IN SAME GPA, BUT MAKE THE SHIELDED/HOODED APPLIC 14 DAYS AFTER SEEDING, AND MAKE THE SAME EVALUATIONS

**IR-4 Residue Trial Plan:** JOINT PROJECT WITH CANADA PMC; NAFTA TRIAL SITES: 5-2 10-6

**Comments:** MFG REQUIRES E/CS DATA PRIOR TO RESIDUE STUDY:09/12; TOLERANCE IS ESTABLISHED, PRE-PLANT USE IS REGISTERED:09/14; E/CS RESEARCH IS ONGOING, BUT NEED MORE POST-TRANSPLANT E/CS DATA BEFORE RESIDUE STUDY:06/17; MFG GAVE OK FOR RESIDUE STUDY AT FUW, SIMILAR TO CABBAGE (10556):09/17; POTENTIAL JOINT PROJECT WITH CANADA:10/17

**NER-EPA Region-FRD**

**NCR-EPA Region-FRD**

**SOR-EPA Region-FRD**

**WSR-EPA Region-FRD**

**CANADA-EPA Region-FRD**

18-GAP01 Culpepper, S.



# 2018 Tentative/Scheduled Studies

## Residue and Efficacy/Crop Safety (E/CS)

(Order by Crop Group, Commodity, Chemical)

Print Date: 2/22/2018

---

<u>PR #</u>	<u>LAB</u>	<u>PRIORITY</u>	<u>STUDY DIRECTOR</u>	<u>CHEMICAL (MFG)</u>	<u>COMMODITY</u>	<u>CROP GROUP</u>
P11591	-	A	DORSCHNER	INSECTICIDE (TBD)	CABBAGE	BRASSICA HEAD AND STEM VEGETABLE GROUP (05-16)

Reason for Need: CABBAGE MAGGOT; NEED NEW ALTERNATIVES TO CHLORPYRIFOS

Use Pattern: (PCR): IN TRAY DRENCH, SEED TREATMENTS, IN-FURROW OR POST-PLANT

E/CS Data Requirements: SUGGESTION IS FOR 2 TRIALS WITH 2 CROPS EACH, ~8 TRTS, POSSIBLY INCLUDING SEED TRT:10/17

E/CS Research Comments:

IR-4 Residue Trial Plan:

Comments: CONSIDER TRIALS ON MULTIPLE CROPS:10/17

NER-EPA Region-FRD

18-NYP03 Gilrein, Dan

NCR-EPA Region-FRD

SOR-EPA Region-FRD

WSR-EPA Region-FRD

18-ORP05 Buckland, Kristine

CANADA-EPA Region-FRD



## 2018 Tentative/Scheduled Studies

### Residue and Efficacy/Crop Safety (E/CS)

*(Order by Crop Group, Commodity, Chemical)*

*Print Date: 2/22/2018*

<b><u>PR #</u></b>	<b><u>LAB</u></b>	<b><u>PRIORITY</u></b>	<b><u>STUDY DIRECTOR</u></b>	<b><u>CHEMICAL (MFG)</u></b>	<b><u>COMMODITY</u></b>	<b><u>CROP GROUP</u></b>
10556	18-FLR01	A	ARSENOVIC	SULFENTRAZONE (FMC)	CABBAGE	BRASSICA HEAD AND STEM VEGETABLE GROUP (05-16)

**Reason for Need:** HOPHORNBEAN COPPERLEAF, PIGWEEDS; NO OTHER PRODUCTS REGISTERED FOR NUTSEDGE CONTROL IN BRASSICA CROPS

**Use Pattern: (PCR):** POST-TRANSPLANT DIRECTED BETWEEN THE ROWS; 2.25-12.0 OZ A/A; 1 SOIL APPLIC; 30-DAY PHI; PER ME-TOO REQUEST, NEED 30-DAY PHI, WHETHER FOR TRANSPLANTED OR DIRECT-SEEDED CROP; PRE-EMERGENCE USE IS COVERED:09/17

**E/CS Data Requirements:**

**E/CS Research Comments:**

**IR-4 Residue Trial Plan:** JOINT PROJECT WITH CANADA PMC; NAFTA TRIAL SITES: 1-2 2-2 5-2 10-2

**Comments:** MFG REQUIRES E/CS DATA PRIOR TO RESIDUE STUDY:09/12; FOR TN, THERE ARE LIMITED TOOLS WITH EFFICACY, AND THIS WOULD INCREASE POTENTIAL FOR ROTATION:06/16; MFG CONFIRMED ONLY RESIDUE DATA NEEDED:06/17; POTENTIAL JOINT PROJECT WITH CANADA:10/17

**NER-EPA Region-FRD**

18-NJ238 Fisher, Jennifer

**NCR-EPA Region-FRD**

18-OH\*265 Horst, Leona

**SOR-EPA Region-FRD**

18-GA\*141 Fraelich, Ben  
(reg 2)  
18-NC214 Batts, Roger B.  
18-SC\*320 Wade, Paul

**WSR-EPA Region-FRD**

18-CA\*28 Benzen, Ms. Sharon D.  
18-CA29 Leach, Nathan  
18-NM253 Hamilton, Cary  
(reg 10)

**CANADA-EPA Region-FRD**

18-NS262 Hanscomb, Darrell D  
18-ON278 Riddle, Geoff  
18-QC312 Cloutier, Dominic



## 2018 Tentative/Scheduled Studies

### Residue and Efficacy/Crop Safety (E/CS)

(Order by Crop Group, Commodity, Chemical)

Print Date: 2/22/2018

<b>PR #</b>	<b>LAB</b>	<b>PRIORITY</b>	<b>STUDY DIRECTOR</b>	<b>CHEMICAL (MFG)</b>	<b>COMMODITY</b>	<b>CROP GROUP</b>
12027	18-YAR02	A	BARNEY	UNICONAZOLE-P (VALENT)	CROP GROUP 05-16 (GH TRANSPLANT)	BRASSICA HEAD AND STEM VEGETABLE GROUP (05-16)

**Reason for Need:** GROWTH REGULATION - WITHOUT THIS, SOME CROP PLANTS WILL BOLT UNDER HOT GH CONDITIONS

**Use Pattern: (PCR):** USE THE SUMAGIC PRODUCT; MAKE UP TO 2 FOLIAR APPLIC OF 2-10 PPM, IN A VOLUME OF 2 QT SOLUTION/100 SQ FT; 7-14 DAY INTERVAL; NO PHI LISTED ON CURRENT LABEL; MFG REQUESTS THE LABELED USE PATTERN FOR FRUITING VEGETABLE TRANSPLANTS BE FOLLOWED (08/16)

**E/CS Data Requirements:**

**E/CS Research Comments:**

**IR-4 Residue Trial Plan:** BROCCOLI: 6 10-4 12; CABBAGE: 1 2 3 5 6 10

**Comments:** REQUEST INCLUDES USE ON VARIOUS GH TRANSPLANTS FOR RETAIL SALE: ROOT/TUBER, LEAFY VEG, BRASSICA, HERBS; THE CURRENT SUMAGIC LABEL INCLUDES ONLY FRUITING VEGETABLES:07/16; MFG SUPPORTS, AND RECOMMENDS THE USE PATTERN CURRENTLY ESTABLISHED FOR FRUITING VEGETABLE TRANSPLANTS:08/16; CONSIDER RESIDUE WORK ON BOTH REP CROPS (CABBAGE AND BROCCOLI) TO COVER THE WHOLE CROP GROUP 5-16:09/16; IR-4 TO PURSUE TOLERANCE WITH CHEMSAC PROPOSAL BASED ON USING DATA TO BE GENERATED ON MUSTARD GREENS (PR# 12026):10/13/16; NEED RESIDUES ON BROCCOLI AND CABBAGE (2 "A" PRIORITIES):09/17

**NER-EPA Region-FRD**

18-MD175 Ross, Marylee  
(cabbage)

**NCR-EPA Region-FRD**

18-MI191 Hausbeck, Dr. Mary K.  
(cabbage)

**SOR-EPA Region-FRD**

18-FL121 Thomas, Darrell  
(cabbage)  
18-NC216 Batts, Roger B.  
(cabbage)  
18-SC\*322 Wade, Paul  
(cabbage)  
18-TX340 Marconi, Cristina  
(broccoli)  
18-TX341 Marconi, Cristina  
(cabbage)

**WSR-EPA Region-FRD**

18-CA\*51 Benzen, Ms. Sharon D.  
(broccoli)  
18-CA52 Ennes, D. (Kearney)  
(broccoli)  
18-CA53 Ennes, D. (Kearney)  
(cabbage)  
18-NM255 Hamilton, Cary  
(broccoli)(reg 10)  
18-OR295 Koskela, Ms. Gina  
(broccoli)  
18-NM349 Hamilton, Cary  
(broccoli)(Reg 10)  
18-OR350 Koskela, Ms. Gina  
(broccoli)(Reg 12)

**CANADA-EPA Region-FRD**



## 2018 Tentative/Scheduled Studies

### Residue and Efficacy/Crop Safety (E/CS)

*(Order by Crop Group, Commodity, Chemical)*

*Print Date: 2/22/2018*

<b><u>PR #</u></b>	<b><u>LAB</u></b>	<b><u>PRIORITY</u></b>	<b><u>STUDY DIRECTOR</u></b>	<b><u>CHEMICAL (MFG)</u></b>	<b><u>COMMODITY</u></b>	<b><u>CROP GROUP</u></b>
12345	18-MIR09	A	LENNON	OXATHIPIPROLIN + MANDIPROPAMID (SYNGEN)	BEAN, LIMA (SUCCULENT)	SUCCULENT/DRIED SHELLED PEA/BEAN SUBGROUPS (06BC)

**Reason for Need:** DOWNY MILDEW AND PHYTOPHTHORA CAPSICI

**Use Pattern: (PCR):** USE THE ORONDIS ULTRA PRODUCT (OXATHIPIPROLIN + MANDIPROPAMID); MAKE FOLIAR APPLIC OF 5.5-8.0 FL OZ/A (INCLUDING AERIAL);  
MAXIMUM ANNUAL RATE 32.0 FL OZ/A

**E/CS Data Requirements:**

**E/CS Research Comments:**

**IR-4 Residue Trial Plan:** 2-4 5 10-2 11; 1 TRIAL IS A DECLINE

**Comments:** THERE ARE TOLERANCES FOR PEA (EDIBLE-PODDED AND SUCCULENT-SHELLED):08/17; SYNGENTA/DUPONT AGREE THIS IS SUPPORTED:09/17; REGISTRANT HAS NOW REQUESTED TO USE A DUAL AI PRODUCT FOR THIS STUDY, OXATHIPIPROLIN + MANDIPROPAMID (FOR RESISTANCE MANAGEMENT AGAINST THE TARGET DISEASES); EACH OF THESE AIS IS CLASSIFIED AS REDUCED RISK:01/18

**NER-EPA Region-FRD**

18-MD179    Ross, Marylee  
18-NJ245    Freiberger, Tom

**NCR-EPA Region-FRD**

18-OH\*274    Horst, Leona  
18-WI383    Heider, Daniel J.

**SOR-EPA Region-FRD**

18-NC221    Batts, Roger B.  
18-SC\*328    Wade, Paul

**WSR-EPA Region-FRD**

18-CA\*82    Benzen, Ms. Sharon D.  
18-CA83    Ennes, D. (Kearney)  
18-CA84    Kyser, Guy  
(decline)  
18-ID163    Meeks, Mr. Will  
18-WA\*366    Harvey, John

**CANADA-EPA Region-FRD**



## 2018 Tentative/Scheduled Studies

### Residue and Efficacy/Crop Safety (E/CS)

*(Order by Crop Group, Commodity, Chemical)*

*Print Date: 2/22/2018*

<b>PR #</b>	<b>LAB</b>	<b>PRIORITY</b>	<b>STUDY DIRECTOR</b>	<b>CHEMICAL (MFG)</b>	<b>COMMODITY</b>	<b>CROP GROUP</b>
11980	17-FLR08	A	SAMOIL	TRIBENURON-METHYL (HSB,NUFARM)	BEAN (DRIED SHELLED)	DRIED SHELLED PEA/BEAN (EXCEPT SOYBEAN) SUBGROUP (06C)

**Reason for Need:** NARROWLEAF HAWKSBEARD, FALSE CHAMOMILE, DANDELION, PRICKLY LETTUCE, MUSTARD SPP.

**Use Pattern: (PCR):** USE THE EXPRESS PRODUCT; MAKE 1 PREPLANT (1-7 DAYS PRIOR TO PLANTING) APPLIC OF 0.25 OZ/A (7 G AI/A) TO EMERGED WEEDS; IF USED ON LIGHT TEXTURED SOILS (SAND AND LOAMY SAND) OR ON HIGH PH SOILS (>7.9), EXTEND TIME TO PLANTING BY 7 MORE DAYS

**E/CS Data Requirements:** CROP SAFETY DATA SHOULD BE COLLECTED DURING RESIDUE TRIAL CONDUCT:09/16

**E/CS Research Comments:**

**IR-4 Residue Trial Plan:** 5-4 7 8 9 10 11 (NO DECLINE TRIAL NEEDED; NO FORAGE OR HAY NEEDED); "RED A" TRIALS NEEDED IN 2018 IN REG. 7 AND 10 (1 TRIAL IN EACH)

**Comments:** KEY EXPORT MARKETS INCLUDE MEXICO, CARRIBEAN, CENTRAL/SOUTH AMERICA, EU; USE IS LABELED IN CANADA, SO CROP TOLERANCE AND YIELD DATA SHOULD BE AVAILABLE:07/16; DUPONT GAVE MFG SUPPORT FOR THIS REQUEST, RESIDUE ONLY, AT THE FUW, PROVIDED THAT CROP SAFETY DATA ARE COLLECTED DURING RESIDUE TRIAL CONDUCT, AND THAT NO FURTHER STUDIES ARE TRIGGERED:09/16

**NER-EPA Region-FRD**

**NCR-EPA Region-FRD**

**SOR-EPA Region-FRD**

**WSR-EPA Region-FRD**

**CANADA-EPA Region-FRD**

18-ND222      Jenks, Dr. Brian  
(reg 7)

18-CA2      Kyser, Guy  
(2018 \$)  
18-CA49      Kyser, Guy  
replacement for .18-CA2  
18-CA\*50      Benzen, Ms. Sharon D.



# 2018 Tentative/Scheduled Studies

## Residue and Efficacy/Crop Safety (E/CS)

(Order by Crop Group, Commodity, Chemical)

Print Date: 2/22/2018

<b>PR #</b>	<b>LAB</b>	<b>PRIORITY</b>	<b>STUDY DIRECTOR</b>	<b>CHEMICAL (MFG)</b>	<b>COMMODITY</b>	<b>CROP GROUP</b>
12261	18-FLR05	A	DORSCHNER	SULFOXAFLO (DOWAGR)	PEA (DRY)	DRIED SHELLD PEA/BEAN (EXCEPT SOYBEAN) SUBGROUP (06C)

**Reason for Need:** PEA APHID, LYGUS BUG

**Use Pattern: (PCR):** USE THE TRANSFORM WG PRODUCT; MAKE UP TO 4 FOLIAR APPLIC OF 0.75-1.0 OZ/A, 14-DAY INTERVAL, 7-DAY PHI; TREAT PER LOCAL ECONOMIC THRESHOLDS; APPLY NO MORE THAN 8.5 OZ/A/YR (0.266 LB AI)

**E/CS Data Requirements:**

**E/CS Research Comments:**

**IR-4 Residue Trial Plan:** 7-3 11/12-2 (SEED, VINES, HAY); 1 TRIAL IS A DECLINE; 2 TRIALS COVERED BY TASC \$\$

**Comments:** REQUEST IS FOR DRY PEAS AND LENTILS; KEY EXPORT MARKETS INCLUDE CHINA, INDIA, EU, SOUTH AMERICA; DOING WORK ON DRY PEA WILL COMPLETE SUBGROUP 6C, AS DRY BEAN HAS A 0.2 PPM TOLERANCE:07/17; MFG SUPPORTS, RESIDUE ONLY:09/17

**NER-EPA Region-FRD**

**NCR-EPA Region-FRD**

**SOR-EPA Region-FRD**

**WSR-EPA Region-FRD**

**CANADA-EPA Region-FRD**

18-ND223 Jenks, Dr. Brian  
(decline)(reg 7)  
18-ND224 Jenks, Dr. Brian  
(TASC \$)(reg 7)  
18-SD331 Clay, Dr. Sharon  
(reg 7)  
18-SD332 Clay, Dr. Sharon  
(reg 7)

18-ID158 Meeks, Mr. Will  
(TASC \$)  
18-ID159 Meeks, Mr. Will  
(reg 11)  
18-WA358 Peng, Wilson  
18-WA\*359 Harvey, John



# 2018 Tentative/Scheduled Studies

## Residue and Efficacy/Crop Safety (E/CS)

(Order by Crop Group, Commodity, Chemical)

Print Date: 2/22/2018

<b><u>PR #</u></b>	<b><u>LAB</u></b>	<b><u>PRIORITY</u></b>	<b><u>STUDY DIRECTOR</u></b>	<b><u>CHEMICAL (MFG)</u></b>	<b><u>COMMODITY</u></b>	<b><u>CROP GROUP</u></b>
12021	17-TBD	A	ARSENOVIC	GLUFOSINATE (BAYER,UPI)	TOMATO	TOMATO SUBGROUP (08-10A)

**Reason for Need:** ANNUAL BROADLEAF WEEDS AND GRASS SPECIES; WILL HELP MANAGE ALS AND GLYPHOSATE RESISTANT WEEDS, AND HELP WITH SEDGES, BOTH PREPLANT AND ROW MIDDLES

**Use Pattern: (PCR):** USE LIBERTY 280 OR RELY 280 PRODUCT; APPLY 0.75 LB AI/A POSTEMERGENCE TO WEEDS, AFTER CROP ESTABLISHMENT OR POST-TRANSPLANTING; NO PHI SPECIFIED; USE A HOODED OR SHIELDED APPLIC; CAN BE USED WITH PLASTICULTURE, OR ROW MIDDLES IN WIDE-ROW CROPS:08/16; FROM GA STAKEHOLDER, USE PATTERNS THAT NEED TO BE COVERED WITH THE RESIDUE PROGRAM INCLUDE: PREPLANT BAREGROUND TRANSPLANTS, PREPLANT AND PRE SEEDED, PREPLANT OVER MULCH WASH OFF PRIOR TO TRANSPLANT, AND ROW MIDDLE HOODED SPRAY:09/16

**E/CS Data Requirements:**

**E/CS Research Comments:** TESTING RELY 280 PRODUCT, WITH THE SAME TREATMENTS IN BARE GROUND AND PLASTIC MULCH CULTURE; COMPARE 0.44, 0.88 AND 1.17 LB AI/A RATES APPLIED BROADCAST TO THE SOIL OR OVER THE MULCH, PRE-TRANSPLANT, WITH THE 0.44 LB AI/A RATE APPLIED TWICE 10-14 DAYS APART STARTING WHEN WEEDS ARE 3-4 INCHES TALL OR 2 WK AFTER TRANSPLANTING, USING A HOODED/SHIELDED BAND TO ROW MIDDLES; ALL RATES ARE APPLIED IN >15 GPA; EVALUATE CROP YIELD AND CROP INJURY; SEE PROTOCOL FOR OTHER APPLIC DETAILS

**IR-4 Residue Trial Plan:** 1 2 3-2 5 10-7, 2 DECLINE TRIALS, 1 PROCESSING TRIAL (PASTE & PUREE)

**Comments:** MFG SUPPORTS:08/16; EPA CAUTION:09/16; MIR IS STORING SAMPLES UNTIL A CONTRACT LAB IS ASSIGNED:09/17

**NER-EPA Region-FRD**

**NCR-EPA Region-FRD**

**SOR-EPA Region-FRD**

**WSR-EPA Region-FRD**

**CANADA-EPA Region-FRD**

18-NM3      Hamilton, Cary  
(Replacement for 17-NM292)(2018 \$)





# 2018 Tentative/Scheduled Studies

## Residue and Efficacy/Crop Safety (E/CS)

(Order by Crop Group, Commodity, Chemical)

Print Date: 2/22/2018

<b>PR #</b>	<b>LAB</b>	<b>PRIORITY</b>	<b>STUDY DIRECTOR</b>	<b>CHEMICAL (MFG)</b>	<b>COMMODITY</b>	<b>CROP GROUP</b>
P12021	-	A	ARSENOVIC	GLUFOSINATE (BAYER,UPI)	TOMATO	TOMATO SUBGROUP (08-10A)

**Reason for Need:** ANNUAL BROADLEAF WEEDS AND GRASS SPECIES; WILL HELP MANAGE ALS AND GLYPHOSATE RESISTANT WEEDS, AND HELP WITH SEDGES, BOTH PREPLANT AND ROW MIDDLES

**Use Pattern: (PCR):** USE LIBERTY 280 OR RELY 280 PRODUCT; APPLY 0.75 LB AI/A POSTEMERGENCE TO WEEDS, AFTER CROP ESTABLISHMENT OR POST-TRANSPLANTING; NO PHI SPECIFIED; USE A HOODED OR SHIELDED APPLIC; CAN BE USED WITH PLASTICULTURE, OR ROW MIDDLES IN WIDE-ROW CROPS:08/16; FROM GA STAKEHOLDER, USE PATTERNS THAT NEED TO BE COVERED WITH THE RESIDUE PROGRAM INCLUDE: PREPLANT BAREGROUND TRANSPLANTS, PREPLANT AND PRE SEEDED, PREPLANT OVER MULCH WASH OFF PRIOR TO TRANSPLANT, AND ROW MIDDLE HOODED SPRAY:09/16

**E/CS Data Requirements:**

**E/CS Research Comments:** TESTING RELY 280 PRODUCT, WITH THE SAME TREATMENTS IN BARE GROUND AND PLASTIC MULCH CULTURE; COMPARE 0.44, 0.88 AND 1.17 LB AI/A RATES APPLIED BROADCAST TO THE SOIL OR OVER THE MULCH, PRE-TRANSPLANT, WITH THE 0.44 LB AI/A RATE APPLIED TWICE 10-14 DAYS APART STARTING WHEN WEEDS ARE 3-4 INCHES TALL OR 2 WK AFTER TRANSPLANTING, USING A HOODED/SHIELDED BAND TO ROW MIDDLES; ALL RATES ARE APPLIED IN >15 GPA; EVALUATE CROP YIELD AND CROP INJURY; SEE PROTOCOL FOR OTHER APPLIC DETAILS

**IR-4 Residue Trial Plan:** 1 2 3-2 5 10-7, 2 DECLINE TRIALS, 1 PROCESSING TRIAL (PASTE & PUREE)

**Comments:** MFG SUPPORTS:08/16; EPA CAUTION:09/16; MIR IS STORING SAMPLES UNTIL A CONTRACT LAB IS ASSIGNED:09/17

**NER-EPA Region-FRD**

18-DEP01 VanGessel, M.  
(DE/NJ ft ID# tbd until researcher assig

**NCR-EPA Region-FRD**

**SOR-EPA Region-FRD**

**WSR-EPA Region-FRD**

18-TBD TBD-WSR  
(CA ft ID# tbd until researcher assigned)(

**CANADA-EPA Region-FRD**



## 2018 Tentative/Scheduled Studies

### Residue and Efficacy/Crop Safety (E/CS)

*(Order by Crop Group, Commodity, Chemical)*

*Print Date: 2/22/2018*

<u>PR #</u>	<u>LAB</u>	<u>PRIORITY</u>	<u>STUDY DIRECTOR</u>	<u>CHEMICAL (MFG)</u>	<u>COMMODITY</u>	<u>CROP GROUP</u>
11894	18-CAR08	A	DORSCHNER	CYCLANILIPROLE (ISK)	TOMATO (GH)	TOMATO SUBGROUP (08-10A)

**Reason for Need:** THRIPS, LEPIDOPTERA

**Use Pattern: (PCR):** USE THE CYCLAPRYN PRODUCT; MAKE FOLIAR APPLIC OF 80-100 G AI/HA, 0-1 DAY PHI

**E/CS Data Requirements:** MFG HAS PERFORMANCE DATA FROM FIELD TOMATO TRIALS, AND IF ACCEPTABLE TO IR-4, NO ADDITIONAL EFFICACY DATA ARE NEEDED:09/16

**E/CS Research Comments:**

**IR-4 Residue Trial Plan:** ANY 4 TRIALS (2 SMALL-FRUITED, 2 LARGE-FRUITED); 1 SMALL-FRUITED TRIAL IS A DECLINE

**Comments:** MFG MAY PROVIDE SOME FINANCIAL GRANT TO OFFSET COSTS:03/16; CANADA PMC HAS GH PEPPER DATA (2017 STUDY) AND WILL HAVE GH CUCUMBER DATA (2018 STUDY):09/17; CANADA TO USE IR-4 GH TOMATO DATA, AND GIVE IR-4 GH PEPPER (11891) AND GH CUCUMBER (11893) DATA, AND 1 EFFICACY TRIAL ON THRIPS:10/17

**NER-EPA Region-FRD**

18-MD173    Ross, Marylee  
(large fruited)

**NCR-EPA Region-FRD**

18-WI374    Chapman, Scott  
(large)

**SOR-EPA Region-FRD**

18-FL119    Thomas, Darrell  
(small fruit)  
18-TX338    Marconi, Cristina  
(large fruit)

**WSR-EPA Region-FRD**

18-CA43    Ennes, D. (Kearney)  
(small fruit)(decline)

**CANADA-EPA Region-FRD**



# 2018 Tentative/Scheduled Studies

## Residue and Efficacy/Crop Safety (E/CS)

(Order by Crop Group, Commodity, Chemical)

Print Date: 2/22/2018

---

<b>PR #</b>	<b>LAB</b>	<b>PRIORITY</b>	<b>STUDY DIRECTOR</b>	<b>CHEMICAL (MFG)</b>	<b>COMMODITY</b>	<b>CROP GROUP</b>
12300	18-CANADA	A	CANADA	NA11630 (TBD)	TOMATO (GH)	TOMATO SUBGROUP (08-10A)

**Reason for Need:** THRIPS, WHITEFLIES, APHIDS

**Use Pattern: (PCR):** MAKE MULTIPLE FOLIAR APPLIC, LOW USE RATE, 7-14 DAY INTERVAL, 0-1 DAY PHI; OTHER DIRECTIONS OF USE PER MFG

**E/CS Data Requirements:** A MINIMUM # OF TRIALS ARE NEEDED:08/17

**E/CS Research Comments:**

**IR-4 Residue Trial Plan:** ANY 4 TRIALS

**Comments:** 2017 FUW "B" PRIORITY, 2017 GLOBAL "A" PRIORITY

**NER-EPA Region-FRD**

**NCR-EPA Region-FRD**

**SOR-EPA Region-FRD**

**WSR-EPA Region-FRD**

**CANADA-EPA Region-FRD**

18-NY393 Palmer, Scott

18-WI396 Heider, Daniel J.

18-FL394 Thomas, Darrell

18-TX395 Marconi, Cristina



# 2018 Tentative/Scheduled Studies

## Residue and Efficacy/Crop Safety (E/CS)

(Order by Crop Group, Commodity, Chemical)

Print Date: 2/22/2018

<u>PR #</u>	<u>LAB</u>	<u>PRIORITY</u>	<u>STUDY DIRECTOR</u>	<u>CHEMICAL (MFG)</u>	<u>COMMODITY</u>	<u>CROP GROUP</u>
11906	18-CAR09	A	LENNON	FLUOPYRAM + PYRIMETHANIL (BAYER)	PEPPER (BELL & NONBELL) (GH)	PEPPER/NON-BELL PEPPER/EGGPLANT SUBGROUPS (08-10BC)

**Reason for Need:** POWDERY MILDEW, BOTRYTIS

**Use Pattern: (PCR):** USE THE LUNA TRANQUILITY PRODUCT; APPLY 100-500 ML PROD/HA, VIA FOLIAR (1000-1500 L/HA SPRAY VOLUME) AND DRENCH, 2 APPLIC 7-12 DAYS APART; 0-1 DAY PHI

**E/CS Data Requirements:**

**E/CS Research Comments:**

**IR-4 Residue Trial Plan:** ANY 4 TRIALS (2 NON-BELL, 2 BELL); 1 NON-BELL TRIAL IS A DECLINE

**Comments:** KEY EXPORT MARKET IS CANADA; THERE IS A PEPPER/EGGPLANT SUBGROUP 8-10B 4.0 PPM TOLERANCE FOR FLUOPYRAM, BUT NO PEPPER TOLERANCE FOR PYRIMETHANIL; THERE ARE OTHER GH USES FOR EACH AI:04/16: MFG IS CONCERNED ABOUT POTENTIAL TRADE IRRITANT SINCE THERE ARE NO CANADIAN PROJECTS TO OBTAIN PYRIMETHANIL REGISTRATION ON GH PEPPER; IF GROWERS ARE OK WITH THIS ISSUE, MFG WOULD SUPPORT THIS USE WITH THE CONDITION THAT A PHYTO WARNING BE PLACED ON THE LABEL SIMILAR TO THE STATEMENT CURRENTLY ON THE SCALA LABEL FOR GH CUCUMBER: 09/16; MFG SUPPORTS AT FUW, RESIDUE ONLY:09/22/16; FLUOPYRAM-EPA CAUTION: 08/17; POTENTIAL JOINT PROJECT W/CANADA:09/17; CANADA TO USE IR-4 DATA:10/17

**NER-EPA Region-FRD**

18-MD174 Ross, Marylee  
(bell)

**NCR-EPA Region-FRD**

18-WI375 Chapman, Scott  
(non-bell)(decline)

**SOR-EPA Region-FRD**

18-FL120 Long, Michael  
(bell)  
18-TX339 Marconi, Cristina  
(non-bell)

**WSR-EPA Region-FRD**

18-CA44 Ennes, D. (Kearney)  
(non-bell)

**CANADA-EPA Region-FRD**



# 2018 Tentative/Scheduled Studies

## Residue and Efficacy/Crop Safety (E/CS)

(Order by Crop Group, Commodity, Chemical)

Print Date: 2/22/2018

---

<u>PR #</u>	<u>LAB</u>	<u>PRIORITY</u>	<u>STUDY DIRECTOR</u>	<u>CHEMICAL (MFG)</u>	<u>COMMODITY</u>	<u>CROP GROUP</u>
12298	18-CANADA	A	CANADA	NA11630 (TBD)	PEPPER (GH)	PEPPER/NON-BELL PEPPER/EGGPLANT SUBGROUPS (08-10BC)

**Reason for Need:** THRIPS, WHITEFLIES, APHIDS

**Use Pattern: (PCR):** MAKE MULTIPLE FOLIAR APPLIC, LOW USE RATE, 7-14 DAY INTERVAL, 0-1 DAY PHI; OTHER DIRECTIONS OF USE PER MFG

**E/CS Data Requirements:** A MINIMUM # OF TRIALS ARE NEEDED:08/17

**E/CS Research Comments:**

**IR-4 Residue Trial Plan:** ANY 4 TRIALS

**Comments:** 2017 FUW "B" PRIORITY, 2017 GLOBAL "A" PRIORITY

**NER-EPA Region-FRD**

18-NY397 Palmer, Scott

**NCR-EPA Region-FRD**

18-WI400 Heider, Daniel J.

**SOR-EPA Region-FRD**

18-FL398 Thomas, Darrell  
18-TX-399 Marconi, Cristina

**WSR-EPA Region-FRD**

**CANADA-EPA Region-FRD**



## 2018 Tentative/Scheduled Studies

### Residue and Efficacy/Crop Safety (E/CS)

*(Order by Crop Group, Commodity, Chemical)*

*Print Date: 2/22/2018*

<b>PR #</b>	<b>LAB</b>	<b>PRIORITY</b>	<b>STUDY DIRECTOR</b>	<b>CHEMICAL (MFG)</b>	<b>COMMODITY</b>	<b>CROP GROUP</b>
P12018	-	A	ARSENOVIC	GLUFOSINATE (BAYER,UPI)	CANTALOUPE	MELON SUBGROUP (09A)

**Reason for Need:** ANNUAL BROADLEAF WEEDS AND GRASS SPECIES; WILL HELP MANAGE ALS AND GLYPHOSATE RESISTANT WEEDS, AND HELP WITH SEDGES, BOTH PREPLANT AND ROW MIDDLES

**Use Pattern: (PCR):** USE LIBERTY 280 OR RELY 280 PRODUCT; APPLY 0.75 LB AI/A POSTEMERGENCE TO WEEDS, AFTER CROP ESTABLISHMENT OR POST-TRANSPLANTING; NO PHI SPECIFIED; USE A HOODED OR SHIELDED APPLIC; CAN BE USED WITH PLASTICULTURE, OR ROW MIDDLES IN WIDE-ROW CROPS:08/16; FROM GA STAKEHOLDER, USE PATTERNS THAT NEED TO BE COVERED WITH THE RESIDUE PROGRAM INCLUDE: PREPLANT BAREGROUND TRANSPLANTS, PREPLANT AND PRE SEEDED, PREPLANT OVER MULCH WASH OFF PRIOR TO TRANSPLANT, AND ROW MIDDLE HOODED SPRAY:09/16

**E/CS Data Requirements:** CANADA PMC CONDUCTED 3 VALUE TRIALS IN 2017

**E/CS Research Comments:** IN 2017 PERF. PROTOCOL, TESTING VARIOUS RATES, PLACEMENT AND TIMINGS IN 4 DIFFERENT TYPES OF PLANTINGS (TRANSPLANTED AND SEEDED LOPES IN BARE GROUND AND PLASTIC MULCH CULTURES); ACCESS THE PROTOCOL ON THE IR-4 WEBSITE FOR FULL DETAILS; COLLECTING CROP INJURY AND YIELD DATA

**IR-4 Residue Trial Plan:** 2 5-2 6 10-4, 1 DECLINE TRIAL (THESE ARE NAFTA TRIAL SITES)

**Comments:** NEED TOLERANCE FOR ENTIRE CROP SUBGROUP 9A TO COVER WATERMELONS; MFG SUPPORTS; CANADA INTEREST:08/16; EPA CAUTION:09/16; MIR IS STORING SAMPLES UNTIL A CONTRACT LAB IS ASSIGNED: 9/17

**NER-EPA Region-FRD**

18-NJP02 Besancon, Thierry  
(DE/NJ ft ID# tbd until researcher assign)

**NCR-EPA Region-FRD**

**SOR-EPA Region-FRD**

**WSR-EPA Region-FRD**

18-TBD TBD-WSR  
(CA ft ID# tbd until researcher assigned)

**CANADA-EPA Region-FRD**



# 2018 Tentative/Scheduled Studies

## Residue and Efficacy/Crop Safety (E/CS)

(Order by Crop Group, Commodity, Chemical)

Print Date: 2/22/2018

<b>PR #</b>	<b>LAB</b>	<b>PRIORITY</b>	<b>STUDY DIRECTOR</b>	<b>CHEMICAL (MFG)</b>	<b>COMMODITY</b>	<b>CROP GROUP</b>
12253	18-TBD	A	HOMA	NMG787 (TBD)	MELON	MELON SUBGROUP (09A)

**Reason for Need:** NEMATODES; NEMATODE MANAGEMENT ALTERNATIVES ARE LIMITED

**Use Pattern: (PCR):** MAKE 1-3 APPLIC TO SOIL PER YEAR, VIA SOIL SPRAY, DRENCH OR DRIP, MAX 500 G AI/HA/YR; OR APPLY BY SEED TREATMENT

**E/CS Data Requirements:**

**E/CS Research Comments:**

**IR-4 Residue Trial Plan:** 2 5 6-2 10-4; 1 TRIAL IS A DECLINE (2 TRIALS COVERED BY FAS \$\$\$)

**Comments:** NEED TO CONDUCT SUFFICIENT TRIALS TO SUPPORT A CODEX MRL; MFG CAN ASSIST BY ANALYZING RESIDUE SAMPLES:08/17; TRIALS TO BE DONE ON CANTALOUPE:09/17

**NER-EPA Region-FRD**

**NCR-EPA Region-FRD**

**SOR-EPA Region-FRD**

**WSR-EPA Region-FRD**

**CANADA-EPA Region-FRD**

18-MI195     Zandstra, Dr. Bernard H.  
 18-OH\*268     Horst, Leona

18-NC219     Batts, Roger B.  
 18-GA\*326     Fraelich, Ben  
 18-TX344     Marconi, Cristina  
 18-TX345     Marconi, Cristina

18-CA64     Kyser, Guy  
 (decline)  
 18-CA65     Leach, Nathan  
 (TASC \$)  
 18-CA66     Skiles, Keri  
 (TASC \$)  
 18-NM256     Hamilton, Cary  
 (reg 10)



# 2018 Tentative/Scheduled Studies

## Residue and Efficacy/Crop Safety (E/CS)

(Order by Crop Group, Commodity, Chemical)

Print Date: 2/22/2018

<u>PR #</u>	<u>LAB</u>	<u>PRIORITY</u>	<u>STUDY DIRECTOR</u>	<u>CHEMICAL (MFG)</u>	<u>COMMODITY</u>	<u>CROP GROUP</u>
12019	17-TBD	A	ARSENOVIC	GLUFOSINATE (BAYER,UPI)	CUCUMBER	SQUASH/CUCUMBER SUBGROUP (09B)

**Reason for Need:** ANNUAL BROADLEAF WEEDS AND GRASS SPECIES; WILL HELP MANAGE ALS AND GLYPHOSATE RESISTANT WEEDS, AND HELP WITH SEDGES, BOTH PREPLANT AND ROW MIDDLES

**Use Pattern: (PCR):** USE LIBERTY 280 OR RELY 280 PRODUCT; APPLY 0.75 LB AI/A POSTEMERGENCE TO WEEDS, AFTER CROP ESTABLISHMENT OR POST-TRANSPLANTING; NO PHI SPECIFIED; USE A HOODED OR SHIELDED APPLIC; CAN BE USED WITH PLASTICULTURE, OR ROW MIDDLES IN WIDE-ROW CROPS:08/16; FROM GA STAKEHOLDER, USE PATTERNS THAT NEED TO BE COVERED WITH THE RESIDUE PROGRAM INCLUDE: PREPLANT BAREGROUND TRANSPLANTS, PREPLANT AND PRE SEEDED, PREPLANT OVER MULCH WASH OFF PRIOR TO TRANSPLANT, AND ROW MIDDLE HOODED SPRAY:09/16

**E/CS Data Requirements:** CANADA PMC CONDUCTED 3 VALUE TRIALS IN 2017

**E/CS Research Comments:** IN 2017 PERF. PROTOCOL, TESTING VARIOUS RATES, PLACEMENT AND TIMINGS IN 4 DIFFERENT TYPES OF PLANTINGS (TRANSPLANTED AND SEEDED CUCUMBERS IN BARE GROUND AND PLASTIC MULCH CULTURES); ACCESS THE PROTOCOL ON THE IR-4 WEBSITE FOR FULL DETAILS; COLLECTING CROP INJURY AND YIELD DATA

**IR-4 Residue Trial Plan:** 2-3 3 5-3 10 (4 PICKLING AND 4 SLICING TRIALS), 1 DECLINE TRIAL (THESE ARE NAFTA TRIAL SITES); NEED A REG. 2 "RED A" TRIAL IN 2018

**Comments:** MFG SUPPORTS; CANADA INTEREST:08/16; EPA CAUTION:09/16; MIR IS STORING SAMPLES UNTIL A CONTRACT LAB IS ASSIGNED: 9/17

NER-EPA Region-FRD

NCR-EPA Region-FRD

SOR-EPA Region-FRD

WSR-EPA Region-FRD

CANADA-EPA Region-FRD

18-GA\*144 Fraelich, Ben

18-SC\*321 Wade, Paul





## 2018 Tentative/Scheduled Studies

### Residue and Efficacy/Crop Safety (E/CS)

*(Order by Crop Group, Commodity, Chemical)*

*Print Date: 2/22/2018*

<u>PR #</u>	<u>LAB</u>	<u>PRIORITY</u>	<u>STUDY DIRECTOR</u>	<u>CHEMICAL (MFG)</u>	<u>COMMODITY</u>	<u>CROP GROUP</u>
12252	18-TBD	A	HOMA	NMG787 (TBD)	CUCUMBER	SQUASH/CUCUMBER SUBGROUP (09B)

**Reason for Need:** NEMATODES; NEMATODE MANAGEMENT ALTERNATIVES ARE LIMITED

**Use Pattern: (PCR):** MAKE 1-3 APPLIC TO SOIL PER YEAR, VIA SOIL SPRAY, DRENCH OR DRIP, MAX 500 G AI/HA/YR; OR APPLY BY SEED TREATMENT

**E/CS Data Requirements:**

**E/CS Research Comments:**

**IR-4 Residue Trial Plan:** 2-3 3 5-2 6 10 (1 TRIAL IS A DECLINE)

**Comments:** NEED TO CONDUCT SUFFICIENT TRIALS TO SUPPORT A CODEX MRL; MFG CAN ASSIST BY ANALYZING RESIDUE SAMPLES:08/17

**NER-EPA Region-FRD**

18-MD176    Ross, Marylee

**NCR-EPA Region-FRD**

18-MI194    Zandstra, Dr. Bernard H.  
18-OH\*267    Horst, Leona

**SOR-EPA Region-FRD**

18-FL131    Long, Michael  
18-GA\*145    Fraelich, Ben  
18-NC218    Batts, Roger B.  
18-SC\*325    Wade, Paul  
18-TX343    Marconi, Cristina

**WSR-EPA Region-FRD**

18-CA63    Kyser, Guy  
(decline)

**CANADA-EPA Region-FRD**



# 2018 Tentative/Scheduled Studies

## Residue and Efficacy/Crop Safety (E/CS)

*(Order by Crop Group, Commodity, Chemical)*

*Print Date: 2/22/2018*

<b>PR #</b>	<b>LAB</b>	<b>PRIORITY</b>	<b>STUDY DIRECTOR</b>	<b>CHEMICAL (MFG)</b>	<b>COMMODITY</b>	<b>CROP GROUP</b>
08857	18-CAR01	A	DORSCHNER	BIFENAZATE (MACDERMID,UPI)	CUCUMBER (GH)	SQUASH/CUCUMBER SUBGROUP (09B)

**Reason for Need:** MITES

**Use Pattern: (PCR):** 0.5 LB A/A; 30-50 GPA; ONE FOLIAR APPLIC EVERY 30 DAYS AS NEEDED; 1-DAY PHI (IR-4 SUGGESTS 2 APPLIC AND A 0-DAY PHI - MFG SUPPORTS THIS:09/16)

**E/CS Data Requirements:**

**E/CS Research Comments:**

**IR-4 Residue Trial Plan:** ANY 4 TRIALS (2 SMALL-FRUITED, 2 LARGE-FRUITED); 1 SMALL-FRUITED TRIAL IS A DECLINE

**Comments:** PR# 07511 (FIELD USE) DOES NOT COVER GREENHOUSE USE; MFG MUST SUBMIT LABEL AMENDMENT:06/06; US TOLERANCE EXISTS FOR FIELD CUCUMBER; LABELED IN CANADA:06/08; NO LONGER MFG OBJECTIVE:06/11; CANADIAN INTEREST (ANY 4 TRIALS):09/13

**NER-EPA Region-FRD**

**NCR-EPA Region-FRD**

**SOR-EPA Region-FRD**

**WSR-EPA Region-FRD**

**CANADA-EPA Region-FRD**

18-WI373 Chapman, Scott  
(large fruit)

18-FL109 Thomas, Darrell  
(large fruit)  
18-NC210 Batts, Roger B.  
(small fruit)(decline)  
18-TX336 Marconi, Cristina  
(small fruit)

18-CA16 Ennes, D. (Kearney)  
(large fruit)



## 2018 Tentative/Scheduled Studies

### Residue and Efficacy/Crop Safety (E/CS)

*(Order by Crop Group, Commodity, Chemical)*

*Print Date: 2/22/2018*

<u>PR #</u>	<u>LAB</u>	<u>PRIORITY</u>	<u>STUDY DIRECTOR</u>	<u>CHEMICAL (MFG)</u>	<u>COMMODITY</u>	<u>CROP GROUP</u>
12254	18-TBD	A	HOMA	NMG787 (TBD)	SQUASH	SQUASH/CUCUMBER SUBGROUP (09B)

**Reason for Need:** NEMATODES; NEMATODE MANAGEMENT ALTERNATIVES ARE LIMITED

**Use Pattern: (PCR):** MAKE 1-3 APPLIC TO SOIL PER YEAR, VIA SOIL SPRAY, DRENCH OR DRIP, MAX 500 G AI/HA/YR; OR APPLY BY SEED TREATMENT

**E/CS Data Requirements:**

**E/CS Research Comments:**

**IR-4 Residue Trial Plan:** 1 2 3 5 10; 1 TRIAL IS A DECLINE

**Comments:** NEED TO CONDUCT SUFFICIENT TRIALS TO SUPPORT A CODEX MRL; MFG CAN ASSIST BY ANALYZING RESIDUE SAMPLES:08/17

**NER-EPA Region-FRD**

18-NJ244 Fisher, Jennifer

**NCR-EPA Region-FRD**

18-MI196 Zandstra, Dr. Bernard H.  
18-OH\*269 Horst, Leona

**SOR-EPA Region-FRD**

18-FL132 Thomas, Darrell  
18-GA\*146 Fraelich, Ben  
18-NC220 Batts, Roger B.  
(decline)  
18-SC\*327 Wade, Paul

**WSR-EPA Region-FRD**

18-NM257 Hamilton, Cary  
(reg 10)

**CANADA-EPA Region-FRD**



## 2018 Tentative/Scheduled Studies

### Residue and Efficacy/Crop Safety (E/CS)

*(Order by Crop Group, Commodity, Chemical)*

*Print Date: 2/22/2018*

<u>PR #</u>	<u>LAB</u>	<u>PRIORITY</u>	<u>STUDY DIRECTOR</u>	<u>CHEMICAL (MFG)</u>	<u>COMMODITY</u>	<u>CROP GROUP</u>
12090	17-CAR01	A	HOMA	FLUOPICOLIDE (VALENT)	ORANGE	ORANGE SUBGROUP (10-10A)

Reason for Need:

Use Pattern: (PCR):

E/CS Data Requirements:

E/CS Research Comments:

IR-4 Residue Trial Plan: 3-8 (REPLACEMENT TRIALS FOR 11021 TRIALS LOST DUE TO UNDER-APPLIC); DO 4 TRIALS IN 2017 (1 DECLINE) & 4 "RED A" TRIALS IN 2018 (1 DECLINE)

Comments: SEE PR# 11021 FOR ALL DETAILS ABOUT THIS REQUEST (NEED, IPM COMMENTS, ETC.); THIS NEW PR# WAS CREATED FOR THIS NEED TO COVER REPLACEMENT "RED A" TRIALS, SINCE 11021 DATA WAS ALREADY SUBMITTED TO EPA:10/16

NER-EPA Region-FRD

NCR-EPA Region-FRD

SOR-EPA Region-FRD

WSR-EPA Region-FRD

CANADA-EPA Region-FRD

18-FL124	Long, Michael
18-FL125	Thomas, Darrell
(decline)	
18-FL126	Thomas, Darrell
18-FL127	Long, Michael



## 2018 Tentative/Scheduled Studies

### Residue and Efficacy/Crop Safety (E/CS)

*(Order by Crop Group, Commodity, Chemical)*

*Print Date: 2/22/2018*

<u>PR #</u> 12091	<u>LAB</u> 17-CAR02	<u>PRIORITY</u> A	<u>STUDY DIRECTOR</u> HOMA	<u>CHEMICAL (MFG)</u> FLUOPICOLIDE (VALENT)	<u>COMMODITY</u> GRAPEFRUIT	<u>CROP GROUP</u> GRAPEFRUIT SUBGROUP (10-10C)
----------------------	------------------------	----------------------	-------------------------------	--	--------------------------------	---

Reason for Need:

Use Pattern: (PCR):

E/CS Data Requirements:

E/CS Research Comments:

IR-4 Residue Trial Plan: 3-3 (REPLACEMENT TRIALS FOR 11022 TRIALS LOST DUE TO UNDER-APPLICATION); DOING 1 TRIAL IN 2017 & 2 "RED A" TRIALS IN 2018

Comments: SEE PR# 11022 FOR ALL DETAILS ABOUT THIS REQUEST (NEED, IPM COMMENTS, ETC.); THIS NEW PR# WAS CREATED FOR THIS NEED TO COVER REPLACEMENT "RED A" TRIALS, SINCE 11022 DATA WAS ALREADY SUBMITTED TO EPA:10/16

NER-EPA Region-FRD

NCR-EPA Region-FRD

SOR-EPA Region-FRD

WSR-EPA Region-FRD

CANADA-EPA Region-FRD

18-FL128	Long, Michael
18-FL129	Thomas, Darrell



## 2018 Tentative/Scheduled Studies

### Residue and Efficacy/Crop Safety (E/CS)

*(Order by Crop Group, Commodity, Chemical)*

Print Date: 2/22/2018

<b>PR #</b>	<b>LAB</b>	<b>PRIORITY</b>	<b>STUDY DIRECTOR</b>	<b>CHEMICAL (MFG)</b>	<b>COMMODITY</b>	<b>CROP GROUP</b>
12259	18-MIR06	A	PIKE	ZETA-CYPERMETHRIN (FMC)	CHERRY	CHERRY SUBGROUP (12-12A)

**Reason for Need:** SPOTTED WING DROSOPHILA - REDUCE PHI FROM 14 TO 3 DAYS TO ALLOW BETTER CROP PROTECTION NEAR HARVEST

**Use Pattern: (PCR):** USE THE MUSTANG MAXX 0.8 EC PRODUCT; MAKE 6 FOLIAR APPLIC OF 0.025 LB AI/A, BY GROUND OR AIR EQUIPMENT, BEGINNING AFTER PETAL FALL, 7-DAY INTERVAL, 3-DAY PHI; USE A MINIMUM 20 GPA FOR CONCENTRATE SPRAY OR 100 GPA FOR DILUTE SPRAY, OR 10 GPA BY AIR; FOR OTHER USE DIRECTIONS FOLLOW THE CURRENT LABEL

**E/CS Data Requirements:**

**E/CS Research Comments:**

**IR-4 Residue Trial Plan:** SOUR: 1 5-4 9; SWEET: 5-2 10-2 11-2 (USING CHERRY, PER SE); 2 TRIALS ARE DECLINES

**Comments:** KEY EXPORT MARKETS INCLUDE CANADA, EU, JAPAN, ASIA; SECTION 18 IS IN PLACE IN MI FOR 2017:06/17; ORIGINAL REQUEST WAS FOR TART CHERRIES, BUT WITH ME-TOO REQUEST FOR SWEET CHERRY, THE COMMODITY WAS CHANGED TO CHERRY, AND COVERS BOTH TART AND SWEET:07/17; EPA CAUTION:08/17; MFG SUPPORTS (AT FUW); WAS A SEC. 18 ON TART CHERRY IN MI IN 2017:09/17

**NER-EPA Region-FRD**

18-NY264 Palmer, Scott  
(sour)

**NCR-EPA Region-FRD**

18-MI197 Van Woerkom, Anthony  
(sour)  
18-MI198 Van Woerkom, Anthony  
(sour)(washed)  
18-MI199 Van Woerkom, Anthony  
(sour)(washed)  
18-MI200 Van Woerkom, Anthony  
(sour)(decline)  
18-MI201 Van Woerkom, Anthony  
(sweet)  
18-MI202 Van Woerkom, Anthony  
(sweet)

**SOR-EPA Region-FRD**

**WSR-EPA Region-FRD**

18-CA67 Skiles, Keri  
(sweet)  
18-CA68 Skiles, Keri  
(sweet)  
18-CA69 Watkins, S.  
(sweet)(decline)  
18-CO103 Oman, Clark (CAT)  
(sour)  
18-OR300 Sturman, Peter  
(sweet)  
18-WA\*356 Harvey, John  
(sweet)  
18-WA357 Peng, Wilson  
(sweet)

**CANADA-EPA Region-FRD**



## 2018 Tentative/Scheduled Studies

### Residue and Efficacy/Crop Safety (E/CS)

*(Order by Crop Group, Commodity, Chemical)*

*Print Date: 2/22/2018*

<b>PR #</b>	<b>LAB</b>	<b>PRIORITY</b>	<b>STUDY DIRECTOR</b>	<b>CHEMICAL (MFG)</b>	<b>COMMODITY</b>	<b>CROP GROUP</b>
P11761	-	A	DORSCHNER	CYFLUMETOFEN (BASF)	PEACH	PEACH SUBGROUP (12-12B)

**Reason for Need:** TWO SPOTTED SPIDER MITE, EUROPEAN RED MITE

**Use Pattern: (PCR):** USE THE NEALTA 200SC PRODUCT; MAKE 2 FOLIAR APPLIC OF 13.7 FL OZ/A IN 50-150 GPA, 7-DAY INTERVAL, 7-DAY PHI

**E/CS Data Requirements:** CANADA DOING 2 PERFORMANCE TRIALS IN 2016

**E/CS Research Comments:** MFG REQUESTS EXAGGERATED RATES TO TEST CROP SAFETY, CONSECUTIVE YEARS TESTING, VARIETY SCREENINGS, ETC.; NEED SUFFICIENT DATA FOR CA:08/15; FOR 2016 CROP SAFETY TESTING, SEE DETAILS OF FOUR TREATMENT REGIMENS IN THE SIGNED PERFORMANCE PROTOCOL; IN 2017 PROTOCOL, TESTING CROP SAFETY ONLY, WITH 2 RATES OF CYFLUMETOFEN (NEALTA), WITH AND WITHOUT NIS ADJUVANT, 2 APPLIC AT A 1-WEEK INTERVAL; COLLECTING CROP INJURY AND CROP YIELD/QUALITY DATA

**IR-4 Residue Trial Plan:** 1 2-3 5 6 10-3 (1 DECLINE); NAFTA SITES: 1 2-2 5-2 10-3

**Comments:** MFG SUPPORTS:08/15; CANADA INTEREST (ZONES 5-4 11):10/15; NY TRIAL LOST IN 2016 WAS REPEATED IN 2016:10/16

**NER-EPA Region-FRD**

**NCR-EPA Region-FRD**

**SOR-EPA Region-FRD**

**WSR-EPA Region-FRD**

**CANADA-EPA Region-FRD**

18-NCP04 Mitchem, Wayne



# 2018 Tentative/Scheduled Studies Residue and Efficacy/Crop Safety (E/CS)

(Order by Crop Group, Commodity, Chemical)

Print Date: 2/22/2018

<b>PR #</b>	<b>LAB</b>	<b>PRIORITY</b>	<b>STUDY DIRECTOR</b>	<b>CHEMICAL (MFG)</b>	<b>COMMODITY</b>	<b>CROP GROUP</b>
10220	18-CAR06	A	HOMA	FLUTIANIL (LANDIS,OATAGRIO)	PEACH	PEACH SUBGROUP (12-12B)

**Reason for Need:** POWDERY MILDEW

**Use Pattern: (PCR):** 6.2-9.4 OZ FORMULATED RATE; FOLIAR VIA AIRBLAST APPLIC IN 100 GPA; 7 APPLIC; 14-DAY RE-TREATMENT INTERVALS

**E/CS Data Requirements:**

**E/CS Research Comments:**

**IR-4 Residue Trial Plan:** 1 2-4 4 5 6 10-4; 2 TRIALS ARE DECLINES

**Comments:** ORIG PCR RECD 08/14/08; HQ HOLD; NEED MFG RESPONSE TO THE INQUIRY OF OTHER POTENTIAL AI PARTNER:07/09; MFG WILL NOT SUPPORT:11/09; HOLD AS THERE ARE NO FOOD TOLERANCES YET; MFG IS RE-EVALUATING:08/16; MFG CONFIRMED THIS REQUEST IS NOW RESEARCHABLE:08/25/16

**NER-EPA Region-FRD**

18-NJ236 Freiburger, Tom  
18-NJ237 Freiburger, Tom  
18-NY263 Palmer, Scott

**NCR-EPA Region-FRD**

18-MI186 Van Woerkom, Anthony

**SOR-EPA Region-FRD**

18-GA\*139 Fraelich, Ben  
18-GA\*140 Fraelich, Ben  
18-NC213 Batts, Roger B.  
(decline)  
18-TX318 Phillips, Mike  
(reg. 6)  
18-LA319 Wright, Denise  
(reg. 4)

**WSR-EPA Region-FRD**

18-CA23 Skiles, Keri  
18-CA24 Skiles, Keri  
18-CA25 Watkins, S.  
(decline)  
18-CA26 Watkins, S.

**CANADA-EPA Region-FRD**





# 2018 Tentative/Scheduled Studies

## Residue and Efficacy/Crop Safety (E/CS)

(Order by Crop Group, Commodity, Chemical)

Print Date: 2/22/2018

<b>PR #</b>	<b>LAB</b>	<b>PRIORITY</b>	<b>STUDY DIRECTOR</b>	<b>CHEMICAL (MFG)</b>	<b>COMMODITY</b>	<b>CROP GROUP</b>
08585	18-MIR03	A	SAMOIL	FLONICAMID (FMC,ISK)	CANEBERRY	CANEBERRY SUBGROUP (13-07A)

**Reason for Need:** STINK BUG, THRIPS

**Use Pattern: (PCR):** 0.067-0.234 LB A/A; 50 GPA (0.13 LB A/A IS MAX RATE MFG SUPPORTS)

**E/CS Data Requirements:**

**E/CS Research Comments:**

**IR-4 Residue Trial Plan:** 1/2 5/6 10 12-3; 1 TRIAL IS A DECLINE; ANY 2 GH TRIALS

**Comments:** PROJECT IS RESEARCHABLE AT 0.089 LB A/A RATE FOR APHIDS/PLANT BUGS; IF HIGHER RATE IS NEEDED FOR STINK BUGS AND THRIPS, PROJECT WILL HAVE TO BE RE-EVALUATED BY MFG:06/09; MFG EFFICACY DATA SHOWS THRIPS SUPPRESSION AT 0.13 LB A/A RATE, SO MFG SUPPORTS RESIDUE WORK:08/15; PER DISCUSSION AT FOOD USE WORKSHOP, TRIALS TO SUPPORT GH USE (PR# 12399) WILL BE INCLUDED WITH THIS PRIORITY "A" PROJECT FOR FIELD USE:09/17

**NER-EPA Region-FRD**

18-NJ232 Freiburger, Tom  
(raspberry)

**NCR-EPA Region-FRD**

18-MI184 Zandstra, Dr. Bernard H.  
(raspberry)

**SOR-EPA Region-FRD**

18-NC209 Batts, Roger B.  
(blackberry)  
18-NC317 Batts, Roger B.  
(gh raspberry)

**WSR-EPA Region-FRD**

18-CA14 Skiles, Keri  
(blackberry)  
18-CA15 TBD-WSR  
(gh raspberry)(check FRD State for ID#)  
18-OR290 Koskela, Ms. Gina  
(blackberry)  
18-OR291 Koskela, Ms. Gina  
(raspberry)  
18-OR292 Koskela, Ms. Gina  
(either decline)

**CANADA-EPA Region-FRD**



# 2018 Tentative/Scheduled Studies

## Residue and Efficacy/Crop Safety (E/CS)

(Order by Crop Group, Commodity, Chemical)

Print Date: 2/22/2018

<b>PR #</b>	<b>LAB</b>	<b>PRIORITY</b>	<b>STUDY DIRECTOR</b>	<b>CHEMICAL (MFG)</b>	<b>COMMODITY</b>	<b>CROP GROUP</b>
P12051	-	+	ARSENOVIC	GLUFOSINATE (BAYER,UPI)	CANEBERRY	CANEBERRY SUBGROUP (13-07A)

**Reason for Need:** ANNUAL, BIENNIAL, AND PERENNIAL BROADLEAF AND GRASS WEEDS THAT ESCAPE WINTER WEED CONTROL STRATEGIES; FOR YEAR-LONG WEED CONTROL PROGRAM; ALTERNATIVE TO PARAQUAT AND GLYPHOSATE; NEEDED FOR RESISTANCE MANAGEMENT

**Use Pattern: (PCR):** USE THE RELY PRODUCT; MAKE 2 POST-EMERGENCE DIRECTED, FOLIAR APPLIC OF 1.5 LB AI/A, 30-DAY INTERVAL, 14-DAY PHI; FIRST APPLIC TO BE MADE BEFORE EMERGENCE OF NEW PRIMOCANES; SECOND APPLIC MUST AVOID DESIRABLE FOLIAGE AND PRIMOCANES

**E/CS Data Requirements:** EVALUATE 8-10 TRTS; APPLY 2 YRS ON THE SAME PLOTS, INCLUDING YR 3 OBSERVATIONS:10/17

**E/CS Research Comments:**

**IR-4 Residue Trial Plan:** 1/2 5/6 10 12-3 (POSSIBLE RESIDUES, DECLINE AND CODEX NEED FOR BLACKBERRY DATA?)

**Comments:** KEY EXPORT MARKETS INCLUDE CANADA, JAPAN, TAIWAN, AUSTRALIA:08/16; IS EPA CAUTION:09/16; MFG SUPPORTS, BUT NEED E/CS DATA BEFORE RESIDUE STUDY; MFG ENCOURAGES JOINT STUDY WITH CANADA:09/16; COULD COVER PR# 05299, OR BE COVERED BY 05299:06/17

**NER-EPA Region-FRD**

**NCR-EPA Region-FRD**

**SOR-EPA Region-FRD**

**WSR-EPA Region-FRD**

**CANADA-EPA Region-FRD**

18-MIP03      Zandstra, Dr. Bernard H.  
(raspberry)

18-WAP02      Miller, Mr. Tim (MT Vern)  
(red raspberry)  
18-CAP08      Smith, Dr. Richard



## 2018 Tentative/Scheduled Studies

### Residue and Efficacy/Crop Safety (E/CS)

*(Order by Crop Group, Commodity, Chemical)*

*Print Date: 2/22/2018*

<u>PR #</u>	<u>LAB</u>	<u>PRIORITY</u>	<u>STUDY DIRECTOR</u>	<u>CHEMICAL (MFG)</u>	<u>COMMODITY</u>	<u>CROP GROUP</u>
08361	18-MIR02	A	BARNEY	ABAMECTIN (SYNGEN)	BLUEBERRY	BUSHBERRY SUBGROUP (13-07B)

**Reason for Need:** BUD MITE AND TWO-SPOTTED SPIDER MITE

**Use Pattern: (PCR):** 20 OZ/A; FOLIAR; POSTHARVEST APPLIC; INCLUDE IN-SEASON APPLIC FOR SPIDER MITES

**E/CS Data Requirements:**

**E/CS Research Comments:**

**IR-4 Residue Trial Plan:** 1 2-3 5-3 12; 1 TRIAL IS A DECLINE

**Comments:** SHOULD BE ABOUT A 10-11 MONTH PHI; USE EXAGGERATED RATE FOR NON-FOOD USE CLASSIFICATION (CHEMSAC PROPOSAL); EPA (HOLD) CAUTION:08/14; EPA CAUTION:08/15; EPA (HOLD) CAUTION:08/16; EPA (HOLD) CAUTION:08/17; NEED SHORT PHI/FOOD USE CLASSIFICATION FOR SPIDER MITE CONTROL:11/17

**NER-EPA Region-FRD**

18-ME104 Collins, Judith  
 18-NJ230 Freiberger, Tom  
 18-NJ231 Freiberger, Tom

**NCR-EPA Region-FRD**

18-MI181 Van Woerkom, Anthony  
 18-MI182 Van Woerkom, Anthony  
 18-MI183 Van Woerkom, Anthony  
 (decline)

**SOR-EPA Region-FRD**

18-GA\*138 Fraelich, Ben  
 18-NC208 Batts, Roger B.

**WSR-EPA Region-FRD**

18-OR289 Koskela, Ms. Gina

**CANADA-EPA Region-FRD**



# 2018 Tentative/Scheduled Studies

## Residue and Efficacy/Crop Safety (E/CS)

(Order by Crop Group, Commodity, Chemical)

Print Date: 2/22/2018

<b>PR #</b>	<b>LAB</b>	<b>PRIORITY</b>	<b>STUDY DIRECTOR</b>	<b>CHEMICAL (MFG)</b>	<b>COMMODITY</b>	<b>CROP GROUP</b>
10614	18-TBD	A	SAMOIL	SULFUR DIOXIDE (TEDMRK)	BLUEBERRY	BUSHBERRY SUBGROUP (13-07B)

**Reason for Need:** BOTRYTIS CINEREA (GREY MOLD) - THERE ARE NO ALTERNATIVES FOR POST-HARVEST CONTROL

**Use Pattern: (PCR):** 100 PPM/HOUR; SINGLE GAS AND/OR SULPHUR DIOXIDE PAD APPLIC IMMEDIATELY AFTER HARVEST AND/OR PACKAGING; SAME SAFETY PRECAUTIONS AS CURRENTLY USED IN TABLE GRAPE INDUSTRY; PER 08/17 REQUESTOR: USE THE FRUIT DOCTOR PRODUCT FROM SNOWDEN ENTERPRISES, INC.; APPLY 200-400 PPM AS POST-HARVEST GAS IN CHAMBER, 0-DAY PHI; AFTER 30 MINUTES EXPOSURE, INITIATE SCRUBBING TO REMOVE RESIDUAL SO2

**E/CS Data Requirements:**

**E/CS Research Comments:**

**IR-4 Residue Trial Plan:** ANY 4 TRIALS (POST HARVEST)

**Comments:** USE IS NEEDED TO PROTECT FLAVOR, EXTEND MARKETING SEASON AND PROLONG POSTHARVEST LIFE; GRAPETEK (PTY) LTD, MANUFACTURER OF UVASYS SULPHUR DIOXIDE GENERATING PADS IN SOUTH AFRICA, RESPONDED THAT THIS REQUEST IS RESEARCHABLE, AND GRAPETEK COULD PROVIDE FUNDING ASSISTANCE FOR THE RESEARCH REQUIRED:07/10; EPA HOLD:07/13; KEY EXPORT MARKETS INCLUDE JAPAN, TAIWAN, SOUTH KOREA; MFG SUPPORTS (PER VERBAL CONFIRMATION AT FUW); TRIALS SHOULD NOT NEED TO BE CONDUCTED USING FRUIT FROM THE 2018 CROP:09/17

**NER-EPA Region-FRD**

**NCR-EPA Region-FRD**

**SOR-EPA Region-FRD**

**WSR-EPA Region-FRD**

**CANADA-EPA Region-FRD**

18-CA35 Ennes, D. (Kearney)  
(special equipment required)  
18-CA36 Ennes, D. (Kearney)  
(special equipment required)  
18-CA37 Ennes, D. (Kearney)  
(special equipment required)  
18-CA38 Ennes, D. (Kearney)  
(special equipment required)



# 2018 Tentative/Scheduled Studies

## Residue and Efficacy/Crop Safety (E/CS)

(Order by Crop Group, Commodity, Chemical)

Print Date: 2/22/2018

<b>PR #</b>	<b>LAB</b>	<b>PRIORITY</b>	<b>STUDY DIRECTOR</b>	<b>CHEMICAL (MFG)</b>	<b>COMMODITY</b>	<b>CROP GROUP</b>
09055	18-CAR04	A	DORSCHNER	INDOXACARB (HSB)	STRAWBERRY	LOW GROWING BERRY SUBGROUP (13-07G)

**Reason for Need:** LEAFROLLER, SAP BEETLE, CLIPPER, PLANT BUG (INCLUDING LYGUS)

**Use Pattern: (PCR):** 0.11 LB AI/A; 2 FOLIAR APPLIC/SEASON; 7-DAY PHI

**E/CS Data Requirements:** SUGGEST NEED FOR GOOD RESULTS (ESPECIALLY FOR LYGUS CONTROL) FROM A MINIMUM 3-4 TRIALS IN LOCATIONS WHERE TARGET PESTS CAN BE E

**E/CS Research Comments:** 2016 PERFORMANCE PROTOCOL INCLUDES 2 RATES OF AVAUNT + NIS ADJUVANT VS A STANDARD, MAX 2 APPLIC NO LESS THAN 7 DAYS APART; PRIMARY TARGET PEST IS TARNISHED PLANT BUG; DATA TO BE COLLECTED INCLUDES PEST CONTROL, CROP INJURY AND CROP YIELD/QUALITY

**IR-4 Residue Trial Plan:** 1 2 3 5 10-3 12; 1 TRIAL IS A DECLINE

**Comments:** EPA CAUTION:08/15; FROM DISCUSSION AT 2015 FUW, CHANGED STATUS FROM NEEDING ONLY RESIDUE DATA; STAKEHOLDERS DECIDED MORE E/CS DATA IS NEEDED BEFORE COMMITTING TO A RESIDUE STUDY; WAS MADE A "+" (H+) FOR THE 2016 PERFORMANCE PROGRAM:09/15; MFG CONFIRMED E/CS DATA REQUIREMENT IS FULFILLED; NOW ONLY NEED RESIDUE DATA:09/15/16; EPA CAUTION:08/17

**NER-EPA Region-FRD**

18-NJ233      Freiburger, Tom  
18-NJ234      Fisher, Jennifer

**NCR-EPA Region-FRD**

18-MI185      Zandstra, Dr. Bernard H.

**SOR-EPA Region-FRD**

18-FL110      Thomas, Darrell  
18-NC211      Batts, Roger B.

**WSR-EPA Region-FRD**

18-CA17      Leach, Nathan  
18-CA18      Skiles, Keri  
(decline)  
18-CA19      Watkins, S.  
18-CA\*20      Benzen, Ms. Sharon D.  
18-OR293      Koskela, Ms. Gina

**CANADA-EPA Region-FRD**



## 2018 Tentative/Scheduled Studies

### Residue and Efficacy/Crop Safety (E/CS)

*(Order by Crop Group, Commodity, Chemical)*

Print Date: 2/22/2018

<u>PR #</u>	<u>LAB</u>	<u>PRIORITY</u>	<u>STUDY DIRECTOR</u>	<u>CHEMICAL (MFG)</u>	<u>COMMODITY</u>	<u>CROP GROUP</u>
12251	18-TBD	A	HOMA	NMG787 (TBD)	STRAWBERRY	LOW GROWING BERRY SUBGROUP (13-07G)

**Reason for Need:** NEMATODES; NEMATODE MANAGEMENT ALTERNATIVES ARE LIMITED

**Use Pattern: (PCR):** MAKE 1-3 APPLIC TO SOIL PER YEAR, VIA SOIL SPRAY, DRENCH OR DRIP, MAX 500 G AI/HA/YR

**E/CS Data Requirements:**

**E/CS Research Comments:**

**IR-4 Residue Trial Plan:** 1 2 3 5 10-3 12; 1 TRIAL IS A DECLINE; FOR JOINT PROJECT WITH PMC, NAFTA TRIAL SITES: 1 3 5-2 10-3 12

**Comments:** NEED TO CONDUCT SUFFICIENT TRIALS TO SUPPORT A CODEX MRL; MFG CAN ASSIST BY ANALYZING RESIDUE SAMPLES:08/17; POTENTIAL JOINT PROJECT WITH CANADA:10/17

**NER-EPA Region-FRD**

18-NJ242 Fisher, Jennifer  
(annual)  
18-NJ243 Freiburger, Tom  
(annual)

**NCR-EPA Region-FRD**

18-MI193 Zandstra, Dr. Bernard H.  
(perennial)  
18-OH\*266 Horst, Leona  
(perennial)

**SOR-EPA Region-FRD**

18-FL130 Long, Michael  
(region 3)(annual)

**WSR-EPA Region-FRD**

18-CA\*59 Benzen, Ms. Sharon D.  
(annual)  
18-CA60 Ennes, D. (Kearney)  
(annual)  
18-CA61 Leach, Nathan  
(annual)  
18-CA62 Watkins, S.  
(decline)(annual)  
18-OR299 Koskela, Ms. Gina  
(reg 12)(perennial)

**CANADA-EPA Region-FRD**

18-ON283 Weber-Henricks, Mary  
(perennial)



# 2018 Tentative/Scheduled Studies

## Residue and Efficacy/Crop Safety (E/CS)

(Order by Crop Group, Commodity, Chemical)

Print Date: 2/22/2018

<u>PR #</u>	<u>LAB</u>	<u>PRIORITY</u>	<u>STUDY DIRECTOR</u>	<u>CHEMICAL (MFG)</u>	<u>COMMODITY</u>	<u>CROP GROUP</u>
P11680	-	A	DORSCHNER	AFIDOPYROPEN (BASF)	STRAWBERRY (GH)	LOW GROWING BERRY SUBGROUP (13-07G)

**Reason for Need:** APHIDS & WHITEFLIES; RESISTANCE MANAGEMENT; NEED ADDITIONAL CONTROL OPTIONS FOR THE GH INDUSTRY

**Use Pattern: (PCR):** FOLIAR APPLIC; OTHER DETAILS TBD BY THE MFG:05/15; MFG REQUESTS THE FOLLOWING: FOLIAR APPLIC AT 7-DAY SPRAY INTERVAL, USING 10-50 G AI/HA, DEPENDING ON TARGET SPECIES; MAX 125 G AI/HA/CROP; PHI TO BE AS SHORT AS POSSIBLE (TBD):09/15

**E/CS Data Requirements:** MFG NEEDS SUFFICIENT DATA FOR REGISTRATION IN CA:08/15; MFG SUPPORTS THE NEED FOR 3 TRIALS WITH CONSISTENT RESULTS:09/15; 1 PMC/CAN

**E/CS Research Comments:** IN 2016 PERFORMANCE PROTOCOL, IN CA AND FL TEST BAS 440 00 I (+ NIS ADJUVANT), AT 20.2 AND 40.4 G AI/A VS A LABELED STANDARD; MAKE 2 FOLIAR BROADCAST APPLIC, WITH THE FIRST WHEN WHITEFLIES REACH A TREATMENT THRESHOLD, FOLLOWED BY A 2ND APPLIC 7 DAYS LATER, IN SUFFICIENT GPA TO PROVIDE THOROUGH COVERAGE; EVALUATE CROP INJURY, PEST CONTROL AND CROP YIELD/QUALITY; IN 2017 PROTOCOL, IN CA TEST THE SAME TREATMENTS USING FOLIAR SPRAYS AND MAKE SIMILAR EVALUATIONS; IN 2018 PROTOCOL, IN GA TEST THE SAME TREATMENTS USING FOLIAR SPRAYS AND MAKE SIMILAR EVALUATIONS

**IR-4 Residue Trial Plan:** ANY 4 TRIALS (1 DECLINE); A LOST BC TRIAL IS BEING REPEATED IN 2017; CA AND TX TRIALS LOST IN 2016 WERE REPLACED WITH 2017 TRIAL #S FOR 2017 FUNDING

**Comments:** ALSO KNOWN BY CODE NAME BAS 440 I:05/15; MFG SUPPORTS FOR BOTH PRODUCTION AND CONSUMER TRANSPLANTS:08/15; MFG MAY ASSIST WITH SAMPLE ANALYSES AND E/CS DATA DEVELOPMENT:09/15; CANADA INTEREST (ANY 3 TRIALS):10/15

NER-EPA Region-FRD

NCR-EPA Region-FRD

SOR-EPA Region-FRD

WSR-EPA Region-FRD

CANADA-EPA Region-FRD

18-GAP02 Riley, D.G.



## 2018 Tentative/Scheduled Studies

### Residue and Efficacy/Crop Safety (E/CS)

*(Order by Crop Group, Commodity, Chemical)*

*Print Date: 2/22/2018*

<b>PR #</b>	<b>LAB</b>	<b>PRIORITY</b>	<b>STUDY DIRECTOR</b>	<b>CHEMICAL (MFG)</b>	<b>COMMODITY</b>	<b>CROP GROUP</b>
11892	18-CAR07	A	SAMOIL	FLUPYRADIFURONE (BAYER)	STRAWBERRY (GH)	LOW GROWING BERRY SUBGROUP (13-07G)

**Reason for Need:** APHIDS, WHITEFLIES; NEED ADDITIONAL CONTROL OPTIONS FOR THE GH INDUSTRY

**Use Pattern: (PCR):** USE SIVANTO PRODUCT (1.67 LB AI/GAL); MAKE 2 FOLIAR APPLIC OF 0.183 LB AI/A IN 20-100 GPA, 7-DAY INTERVAL, 0-DAY PHI

**E/CS Data Requirements:** MFG REQUESTS EFFICACY AND CROP SAFETY DATA:09/17; PMC CANADA DOING 1 PERFORMANCE TRIAL IN 2018:01/18

**E/CS Research Comments:**

**IR-4 Residue Trial Plan:** ANY 4 TRIALS; 1 TRIAL IS A DECLINE; FOR JOINT PROJECT WITH PMC, NAFTA TRIAL SITES: ANY 3 TRIALS

**Comments:** IS LABELED ON FIELD-GROWN STRAWBERRY, WITH A 0-DAY PHI; CAN EXPLORE COVERING GH USE WITH EXISTING TOLERANCE:03/16; MFG SUPPORTS:05/16; MFG MAY NOT SUPPORT USE IN ALL PARTS OF THE COUNTRY DUE TO PEST RESISTANCE CONCERNS:05/17; POTENTIAL JOINT PROJECT WITH CANADA:10/17; RESEARCHABLE, RESIDUE & E/CS DATA NEEDED

**NER-EPA Region-FRD**

18-MD172    Ross, Marylee  
(decline)

**NCR-EPA Region-FRD**

**SOR-EPA Region-FRD**

18-FL118    Long, Michael

**WSR-EPA Region-FRD**

18-CA42    Ennes, D. (Kearney)

**CANADA-EPA Region-FRD**

18-ON280    Riddle, Geoff





## 2018 Tentative/Scheduled Studies Residue and Efficacy/Crop Safety (E/CS)

(Order by Crop Group, Commodity, Chemical)

Print Date: 2/22/2018

<b><u>PR #</u></b>	<b><u>LAB</u></b>	<b><u>PRIORITY</u></b>	<b><u>STUDY DIRECTOR</u></b>	<b><u>CHEMICAL (MFG)</u></b>	<b><u>COMMODITY</u></b>	<b><u>CROP GROUP</u></b>
11937	18-FLR07	A	LENNON	CYPRODINIL + FLUDIOXONIL (SYNGEN)	CRANBERRY	LOW GROWING BERRY SUBGROUP, EXCEPT STRAWBERRY (13-07H)

**Reason for Need:** FRUIT ROT, PHOMOPSIS SPP., COLLETOTRICHUM SPP., ALTERNARIA SPP.

**Use Pattern: (PCR):** USE THE SWITCH PRODUCT; MAKE UP TO 4 FOLIAR APPLIC (VIA GROUND, CHEMIGATION OR AIR) OF 6.9-8.75 OZ/A, 7-10 DAY INTERVALS BEGINNING AT OR BEFORE BLOOM, 30-DAY PHI

**E/CS Data Requirements:**

**E/CS Research Comments:**

**IR-4 Residue Trial Plan:** 1-2 5-2 12; 1 TRIAL IS A DECLINE

**Comments:** KEY EXPORT MARKETS INCLUDE EU, CANADA, CODEX; SUBGROUP 13-07G TOLERANCES FOR BOTH AIS EXCLUDE CRANBERRY DUE TO THE 0-DAY PHI, AND FOR CRANBERRY THE PHI NEEDS TO BE MUCH LONGER:06/16; MFG SUPPORTS ONLY IF LABEL LANGUAGE CAN EXCLUDE DIRECT APPLICATIONS TO WATER:07/16; FLUDIOXONIL IS AN EPA CAUTION:08/16; CYPRODINIL & FLUDIOXONIL-EPA GREENS:08/17

**NER-EPA Region-FRD**

18-MA171    Sylvia, Marty  
18-NJ241    Freiburger, Tom

**NCR-EPA Region-FRD**

18-WI376    Heider, Daniel J.  
18-WI377    Heider, Daniel J.  
(decline)

**SOR-EPA Region-FRD**

**WSR-EPA Region-FRD**

18-OR294    Sturman, Peter

**CANADA-EPA Region-FRD**



## 2018 Tentative/Scheduled Studies

### Residue and Efficacy/Crop Safety (E/CS)

(Order by Crop Group, Commodity, Chemical)

Print Date: 2/22/2018

PR #	LAB	PRIORITY	STUDY DIRECTOR	CHEMICAL (MFG)	COMMODITY	CROP GROUP
12024	18-TBD	A	ARSENOVIC	PROHEXADIONE CALCIUM (BASF,FINEAMA)	CORN (ALFALFA)	CEREAL GRAINS AND CEREAL GRAINS FORAGE/FODDER/STRAW GROUPS (15-16)

**Reason for Need:** PROHEXADIONE IS USED TO ENHANCE ESTABLISHMENT AND SUBSEQUENT YIELD OF ALFALFA INTERSEEDED UNDER CORN; IT IS ANTICIPATED THAT THIS WOULD BE USED ON ABOUT 400,000 ACRES PER YEAR IN THE NORTHERN US WHERE ALFALFA CAN'T BE SUCCESSFULLY ESTABLISHED IN THE FALL AFTER SILAGE CORN HARVEST; MUCH OF THIS CROPLAND NEEDS IMPROVED MANAGEMENT PRACTICES TO REDUCE RUNOFF OF SOIL AND NUTRIENTS AND LEACHING OF NITRATE, AND INTERSEEDED ALFALFA WOULD SERVE AS A DUAL-PURPOSE CROP FOR PROVIDING GROUND COVER DURING SILAGE CORN PRODUCTION AND FORAGE DURING SUBSEQUENT GROWING SEASONS; RECENT STUDIES DEMONSTRATE THAT PROPERLY TIMED PROHEXADIONE ON ALFALFA SEEDLINGS INCREASED PLANT SURVIVAL BY UP TO 400%, GREATLY IMPROVING THE ESTABLISHMENT OF INTERSEEDED ALFALFA IN SILAGE CORN; OTHER STUDIES SHOWED THAT INTERSEEDED ALFALFA REDUCED FALL AND SPRING RUNOFF OF PHOSPHORUS BY 60% AND SOIL EROSION BY 80% COMPARED TO CROPLAND CONTAINING ONLY SILAGE CORN RESIDUES AND WEEDS; ALFALFA IS ALSO KNOWN TO BE HIGHLY EFFECTIVE FOR REDUCING NITRATE LEACHING INTO GROUNDWATER; THESE ARE ALL PPOWERFUL INCENTIVES FOR REGISTERING THIS USE

**Use Pattern: (PCR):** ORIGINAL REQUEST: USE THE "APOGEE" PRODUCT (27.5% WETTABLE GRANULES); MAKE 1 APPLIC OF 6.5-26 OZ PRODUCT/A (WITH POTENTIAL OF UP TO 52 OZ/A) IN A FOLIAR SPRAY DIRECTED TO 6-12 INCH TALL ALFALFA (WITH STEMS RAPIDLY EXPANDING) USING DROP NOZZLES TO MINIMIZE CONTACT WITH CORN PLANTS; NO PHI SPECIFIED; APPLY IN 15-30 GPA, ADDING AMMONIUM SULFATE (2.5 LB/100 GAL) AND CROP OIL CONCENTRATE (1 QT/A) TO MAXIMIZE UPTAKE, AND CITRIC ACID (0.5 G/100 ML) TO ENSURE PH OF SOLUTION IS ADEQUATE FOR ABSORPTION:07/16; FOR FINE AMERICAS PRODUCT "KUDOS" ADJUST USE PATTERN IF NEEDED FOR THIS 27.5 WDG FORMULATION:01/17; CAN INTERSEED THE 2 CROPS IN THE SAME PLOTS:09/17

**E/CS Data Requirements:**

**E/CS Research Comments:**

**IR-4 Residue Trial Plan:** 2-YR STUDY, WI ONLY; ALFALFA/CORN MIXTURE: 3 TRIALS IN 2018, SILAGE

**Comments:** ALFALFA SURVIVAL IS LIMITED UNDER THE COMPETITION OF THE CORN SILAGE CROP; PROHEXADIONE HAS BEEN SHOWN TO INCREASE FIRST YEAR ALFALFA YIELDS BY 12% AND FALL STAND DENSITIES BY 37-130% COMPARED TO UNTREATED INTERSEEDED CONTROLS:08/16; BASF DOES NOT SUPPORT THIS USE WITH PRODUCT "APOGEE":09/16; REQUEST BEING CONSIDERED BY FINE AMERICAS FOR THEIR PRODUCT "KUDOS":12/16; MFG SUPPORTS, ONLY RESIDUE DATA NEEDED (ON ALFALFA AND CORN); 2-YEAR, REGIONAL (STATE - WI) TOLERANCE:09/17

**NER-EPA Region-FRD**

**NCR-EPA Region-FRD**

**SOR-EPA Region-FRD**

**WSR-EPA Region-FRD**

**CANADA-EPA Region-FRD**

18-WI378 Heider, Daniel J.  
(corn)  
18-WI379 Heider, Daniel J.  
(corn)  
18-WI380 Heider, Daniel J.  
(corn)



# 2018 Tentative/Scheduled Studies

## Residue and Efficacy/Crop Safety (E/CS)

(Order by Crop Group, Commodity, Chemical)

Print Date: 2/22/2018

<b>PR #</b>	<b>LAB</b>	<b>PRIORITY</b>	<b>STUDY DIRECTOR</b>	<b>CHEMICAL (MFG)</b>	<b>COMMODITY</b>	<b>CROP GROUP</b>
11979	17-CAR07	A	SAMOIL	METHOXYFENOZIDE (DOWAGR)	RICE	CEREAL GRAINS AND CEREAL GRAINS FORAGE/FODDER/STRAW GROUPS (15-16)

**Reason for Need:** ARMYWORM (TRUE & WESTERN YELLOWSTRIPED); ARMYWORM IS USUALLY CYCLICAL, NEEDING CONTROL ABOUT EVERY 5 YEARS; DROUGHT AND INSECTICIDE RESISTANCE HAVE CAUSED SIGNIFICANT, RECURRING OUTBREAKS; A DIFFERENT MODE OF ACTION ALTERNATIVE TOOL IS NEEDED FOR ARMYWORM CONTROL:07/16

**Use Pattern: (PCR):** USE THE INTREPID 2F PRODUCT; MAKE 2 FOLIAR APPLIC OF 0.12-0.16 LB AI/A, 10-DAY INTERVAL, 14-DAY PHI; BEGIN APPLIC WHEN THRESHOLD LEVEL OF FEEDING DAMAGE OCCURS

**E/CS Data Requirements:**

**E/CS Research Comments:**

**IR-4 Residue Trial Plan:** 3 TRIALS IN 10, GRAIN AND STRAW (OVER 2 YEARS FOR REGIONAL TOLERANCE), 1 DECLINE TRIAL, 1 PROCESSING TRIAL (POLISHED RICE, HULLS, BRAN); PER 10/17-18/16 EMAILS, CODEX REQUIRES MIN 8 TRIALS, SO 4 TRIALS IN 2017, THEN 4 MORE IN 2018

**Comments:** KEY EXPORT MARKET IS JAPAN (ABOUT 50% IS EXPORTED, AND HALF OF THAT GOES TO JAPAN); REPRESENTS A POTENTIAL SECTION 18 NEED; IR-4 SUGGESTS THIS IS A PROJECT FOR CA RICE ONLY (SUSHI RICE):07/16; MFG SUPPORTS AT FUW, RESIDUE ONLY:09/16; EPA RICE GRAIN TOLERANCE IS TREATED THE SAME AS THE JAPANESE BROWN RICE MRL - HARMONIZATION OF THE MRL IS NEEDED:10/16; CA CRISIS EXEMPTIONS WERE APPROVED IN 2015 AND 2016, AND A SPECIFIC EMERGENCY SEC. 18 EXEMPTION FOR 2017 FOR ARMYWORM CONTROL, SUPPORTED BY IR-4'S RESIDUE STUDY DEMONSTRATING PROGRESS TOWARD REGISTRATION:07/17

**NER-EPA Region-FRD**

**NCR-EPA Region-FRD**

**SOR-EPA Region-FRD**

**WSR-EPA Region-FRD**

**CANADA-EPA Region-FRD**

18-CA45	Watkins, S.
18-CA46	Watkins, S.
18-CA47	Watkins, S.
18-CA48	Watkins, S.



# 2018 Tentative/Scheduled Studies

## Residue and Efficacy/Crop Safety (E/CS)

(Order by Crop Group, Commodity, Chemical)

Print Date: 2/22/2018

<b><u>PR #</u></b>	<b><u>LAB</u></b>	<b><u>PRIORITY</u></b>	<b><u>STUDY DIRECTOR</u></b>	<b><u>CHEMICAL (MFG)</u></b>	<b><u>COMMODITY</u></b>	<b><u>CROP GROUP</u></b>
12403	18-MIR13	A	DORSCHNER	CYANTRANILIPROLE (HGW86) (HSB)	BASIL	HERB SUBGROUP (19A)

**Reason for Need:** SEE REASON FOR THIS PR# IN THE COMMENTS, AND THE NEED ON INDIAN CURRY LEAVES FOR ASIAN CITRUS PSYLLID IN PR# 12390

**Use Pattern: (PCR):** FROM REQUEST 12390 FOR INDIAN CURRY LEAVES: USE EXIREL OR VERIMARK PRODUCT; MAKE SOIL APPLIC OF 0.261-0.391 LB AI/A (MFG CONFIRMED THIS RATE RANGE IS FOR VERIMARK) VIA CHEMIGATION, 30-DAY INTERVAL, 1-DAY PHI; APPLY TO FIRST THIRD OF THE IRRIGATION CYCLE; DO NOT IRRIGATE FOR 24 HR AFTER APPLIC; PER MFG, BOTH EXIREL AND VERIMARK ARE EFFICACIOUS, BUT EXIREL (FOLIAR) HAS BEEN THE BEST OPTION, BASED ON USE RATES AND PRICE; FOR EXIREL, RESEARCHER INDICATES RATE RANGE IS 0.088-0.133 LB AI/A IN 100-150 GPA, WITH THE 0.133 LB AI RATE CONSIDERED BEST IN ORDER TO ENSURE EFFICACY:10/17

**E/CS Data Requirements:**

**E/CS Research Comments:**

**IR-4 Residue Trial Plan:** ANY 4 (1 DECLINE, ON FRESH); FRESH AND DRY

**Comments:** PROJECT ADDED FOR THIS FUTURE REP CROP FOR CROP GROUP 25 (HERBS), WHICH WILL PROVIDE A TOLERANCE THAT WILL COVER USE ON INDIAN CURRY LEAVES (SEE PR# 12390):10/17

**NER-EPA Region-FRD**

18-NJ250 Fisher, Jennifer

**NCR-EPA Region-FRD**

18-OH\*276 Horst, Leona  
18-WI391 Chapman, Scott

**SOR-EPA Region-FRD**

18-SC\*330 Wade, Paul

**WSR-EPA Region-FRD**

18-CA97 Leach, Nathan  
18-WA\*372 Harvey, John  
(decline)

**CANADA-EPA Region-FRD**



# 2018 Tentative/Scheduled Studies Residue and Efficacy/Crop Safety (E/CS)

(Order by Crop Group, Commodity, Chemical)

Print Date: 2/22/2018

<u>PR #</u>	<u>LAB</u>	<u>PRIORITY</u>	<u>STUDY DIRECTOR</u>	<u>CHEMICAL (MFG)</u>	<u>COMMODITY</u>	<u>CROP GROUP</u>
12294	18-CAR11	A	SAMOIL	FLUPYRADIFURONE (BAYER)	BASIL (GH)	HERB SUBGROUP (19A)

**Reason for Need:** APHIDS AND WHITEFLIES

**Use Pattern: (PCR):** USE THE ALTUS PRODUCT; MAKE 1 FOLIAR APPLIC OF 7-12 (FURTHER DOSAGE INFO NOT DEFINED) PER MONTH; 7-DAY MINIMUM SPRAY INTERVAL, 0-1 DAY PHI; FOLLOW THE LABEL FOR LETTUCE; BASED ON THE LEAF LETTUCE USE PATTERN, MAKE 2 FOLIAR APPLIC OF 14 FL OZ PRODUCT/A, 7-DAY INTERVAL, 1-DAY PHI; PLUS MAKE 1 DRENCH APPLIC OF 28 FL OZ PRODUCT/A, 1-DAY PHI (RECOMMEND MULTIPLE HARVESTS TO CAPTURE PEAK RESIDUES)

**E/CS Data Requirements:** MFG REQUESTS EFFICACY AND CROP SAFETY DATA; 3 TRIALS ANYWHERE; HIGH RATE, LOW RATE, STANDARD, CONTROL; INCLUDE CHEMIGATION AND F

**E/CS Research Comments:**

**IR-4 Residue Trial Plan:** ANY 4 TRIALS (FRESH ONLY); 1 TRIAL IS A DECLINE

**Comments:** IR-4 SUGGESTS INCLUDING WITH THE FOLIAR USE REQUEST A SOIL USE:07/17; MFG SUPPORTS:09/17

**NER-EPA Region-FRD**

18-MD178 Ross, Marylee  
(decline)

**NCR-EPA Region-FRD**

18-OH\*272 Horst, Leona  
18-WI381 Chapman, Scott

**SOR-EPA Region-FRD**

**WSR-EPA Region-FRD**

18-CA77 Ennes, D. (Kearney)

**CANADA-EPA Region-FRD**



# 2018 Tentative/Scheduled Studies Residue and Efficacy/Crop Safety (E/CS)

(Order by Crop Group, Commodity, Chemical)

Print Date: 2/22/2018

<b><u>PR #</u></b>	<b><u>LAB</u></b>	<b><u>PRIORITY</u></b>	<b><u>STUDY DIRECTOR</u></b>	<b><u>CHEMICAL (MFG)</u></b>	<b><u>COMMODITY</u></b>	<b><u>CROP GROUP</u></b>
P12294	-	A	DORSCHNER	FLUPYRADIFURONE (BAYER)	BASIL (GH)	HERB SUBGROUP (19A)

**Reason for Need:** APHIDS AND WHITEFLIES

**Use Pattern: (PCR):** USE THE ALTUS PRODUCT; MAKE 1 FOLIAR APPLIC OF 7-12 (FURTHER DOSAGE INFO NOT DEFINED) PER MONTH; 7-DAY MINIMUM SPRAY INTERVAL, 0-1 DAY PHI; FOLLOW THE LABEL FOR LETTUCE; BASED ON THE LEAF LETTUCE USE PATTERN, MAKE 2 FOLIAR APPLIC OF 14 FL OZ PRODUCT/A, 7-DAY INTERVAL, 1-DAY PHI; PLUS MAKE 1 DRENCH APPLIC OF 28 FL OZ PRODUCT/A, 1-DAY PHI (RECOMMEND MULTIPLE HARVESTS TO CAPTURE PEAK RESIDUES)

**E/CS Data Requirements:** MFG REQUESTS EFFICACY AND CROP SAFETY DATA; 3 TRIALS ANYWHERE; HIGH RATE, LOW RATE, STANDARD, CONTROL; INCLUDE CHEMIGATION AND F

**E/CS Research Comments:**

**IR-4 Residue Trial Plan:** ANY 4 TRIALS (FRESH ONLY); 1 TRIAL IS A DECLINE

**Comments:** IR-4 SUGGESTS INCLUDING WITH THE FOLIAR USE REQUEST A SOIL USE:07/17; MFG SUPPORTS:09/17

**NER-EPA Region-FRD**

**NCR-EPA Region-FRD**

**SOR-EPA Region-FRD**

**WSR-EPA Region-FRD**

**CANADA-EPA Region-FRD**

18-WIP03 Chapman, Scott

18-GAP04 Riley, D.G.



# 2018 Tentative/Scheduled Studies

## Residue and Efficacy/Crop Safety (E/CS)

(Order by Crop Group, Commodity, Chemical)

Print Date: 2/22/2018

<b><u>PR #</u></b>	<b><u>LAB</u></b>	<b><u>PRIORITY</u></b>	<b><u>STUDY DIRECTOR</u></b>	<b><u>CHEMICAL (MFG)</u></b>	<b><u>COMMODITY</u></b>	<b><u>CROP GROUP</u></b>
12401	18-MIR11	A	DORSCHNER	CYANTRANILIPROLE (HGW86) (HSB)	DILL	HERB SUBGROUP (19A)

**Reason for Need:** SEE REASON FOR THIS PR# IN THE COMMENTS, AND THE NEED ON KAFFIR LIME CITRUS LEAVES FOR ASIAN CITRUS PSYLLID IN PR# 12389

**Use Pattern: (PCR):** FROM REQUEST 12389 FOR KAFFIR LIME CITRUS LEAVES: USE EXIREL OR VERIMARK PRODUCT; MAKE SOIL APPLIC OF 0.261-0.391 LB AI/A (MFG CONFIRMED THIS RATE RANGE IS FOR VERIMARK) VIA CHEMIGATION, 30-DAY INTERVAL, 1-DAY PHI; APPLY TO FIRST THIRD OF THE IRRIGATION CYCLE; DO NOT IRRIGATE FOR 24 HR AFTER APPLIC; PER MFG, BOTH EXIREL AND VERIMARK ARE EFFICACIOUS, BUT EXIREL (FOLIAR) HAS BEEN THE BEST OPTION, BASED ON USE RATES AND PRICE; FOR EXIREL, RESEARCHER INDICATES RATE RANGE IS 0.088-0.133 LB AI/A IN 100-150 GPA, WITH THE 0.133 LB AI RATE CONSIDERED BEST IN ORDER TO ENSURE EFFICACY:10/17

**E/CS Data Requirements:**

**E/CS Research Comments:**

**IR-4 Residue Trial Plan:** ANY 4 TRIALS (1 DECLINE TRIAL); SEED, FRESH AND DRIED LEAVES, OIL (2 SITES ONLY)

**Comments:** PROJECT ADDED FOR THIS FUTURE REP CROP FOR CROP GROUP 26 (SPICES), WHICH WILL PROVIDE A TOLERANCE THAT WILL COVER USE ON KAFFIR LIME CITRUS LEAVES (SEE PR# 12389):10/17

**NER-EPA Region-FRD**

18-NJ249 Fisher, Jennifer

**NCR-EPA Region-FRD**

18-WI388 Chapman, Scott  
(oil)

**SOR-EPA Region-FRD**

18-FL136 Long, Michael

**WSR-EPA Region-FRD**

18-CA96 Ennes, D. (Kearney)  
18-ID167 Meeks, Mr. Will  
(decline)  
18-WA\*370 Harvey, John  
(oil)

**CANADA-EPA Region-FRD**



# 2018 Tentative/Scheduled Studies

## Residue and Efficacy/Crop Safety (E/CS)

(Order by Crop Group, Commodity, Chemical)

Print Date: 2/22/2018

---

<u>PR #</u>	<u>LAB</u>	<u>PRIORITY</u>	<u>STUDY DIRECTOR</u>	<u>CHEMICAL (MFG)</u>	<u>COMMODITY</u>	<u>CROP GROUP</u>
P12349	-	+	HOMA	FLUTIANIL (LANDIS,OATAGRIO)	ROSEMARY	HERB SUBGROUP (19A)

**Reason for Need:** POWDERY MILDEW - IS NOT BEING CONTROLLED WITH REGISTERED FUNGICIDES

**Use Pattern: (PCR):** MAKE FOLIAR APPLIC OF 0.04 LB AI/A, MINIMUM 7-DAY INTERVAL, MAXIMUM 3 APPLIC/CROP; MFG CONSIDERING REGISTERING A LOWER APPLIC RATE THAN REQUESTED

**E/CS Data Requirements:**

**E/CS Research Comments:**

**IR-4 Residue Trial Plan:**

**Comments:** REQUEST IS FOR USE ON FIELD AND GH-GROWN ROSEMARY; COULD BE COVERED IF DATA IS COLLECTED ON REP CROP BASIL; MFG INDICATES POSSIBLE ASSISTANCE WITH ANALYSIS OF RESIDUE SAMPLES:08/17; MFG CHECKING ON NEED FOR E/CS DATA FIRST:09/17

NER-EPA Region-FRD

NCR-EPA Region-FRD

SOR-EPA Region-FRD

WSR-EPA Region-FRD

CANADA-EPA Region-FRD

18-NYP04     Catlin, Nora  
(GH)





## 2018 Tentative/Scheduled Studies

### Residue and Efficacy/Crop Safety (E/CS)

*(Order by Crop Group, Commodity, Chemical)*

*Print Date: 2/22/2018*

<b><u>PR #</u></b>	<b><u>LAB</u></b>	<b><u>PRIORITY</u></b>	<b><u>STUDY DIRECTOR</u></b>	<b><u>CHEMICAL (MFG)</u></b>	<b><u>COMMODITY</u></b>	<b><u>CROP GROUP</u></b>
P12343	-	+	BARNEY	PENDIMETHALIN (BASF,UPI)	ROSEMARY	HERB SUBGROUP (19A)

**Reason for Need:** ANNUAL BROADLEAF AND GRASS WEEDS

**Use Pattern: (PCR):** USE PROWL H2O PRODUCT; MAKE 2-3 APPLIC OF 1-4 PT/A VIA PIVOT IRRIGATION, DIRECTED SPRAY OR COATED FERTILIZER; OTHER USE DIRECTIONS PER LABEL FOR SIMILAR CROPS

**E/CS Data Requirements:** MFG REQUESTS A 2-3 TRIAL CROP SAFETY PROGRAM, AT 1X AND 2X RATES ON ESTABLISHED ROSEMARY, WITH PROWL H2O APPLIED PREEMERGENCE A

**E/CS Research Comments:**

**IR-4 Residue Trial Plan:**

**Comments:** FOR USE IN TRANSPLANTED AND ESTABLISHED PERENNIAL FIELDS; A VERY OLD REQUEST (02476) WAS NOT SUPPORTED AT THAT TIME; THIS CAN POTENTIALLY BE PURSUED WITH A CHEMSAC PROPOSAL, USING THE SAME RATIONALE AS USED FOR PENDIMETHALIN ON MONARDA (PR# 11726); MFG SUPPORTS THIS USE, BUT CROP SAFETY DATA IS NEEDED BEFORE APPROVAL FOR RESIDUE WORK/LABELLING (= "POTENTIAL"), BASED ON THE PRESENCE OF ROSEMARY (WHEN GROWN AS A PERENNIAL ORNAMENTAL PLANT) ALREADY ON THE PENDIMETHALIN LABEL (PENDULUM AQUACAP):08/17

**NER-EPA Region-FRD**

**NCR-EPA Region-FRD**

**SOR-EPA Region-FRD**

**WSR-EPA Region-FRD**

**CANADA-EPA Region-FRD**

18-NMP01 Baker, John  
 Kemin researcher/ft in Hobbs, NM  
 18-TXP01 Baker, John  
 Kemin researcher/ft in Midkiff, TX  
 18-TXP02 Baker, John  
 Kemin researcher/ft in Poteet, TX



## 2018 Tentative/Scheduled Studies

### Residue and Efficacy/Crop Safety (E/CS)

*(Order by Crop Group, Commodity, Chemical)*

*Print Date: 2/22/2018*

<b><u>PR #</u></b>	<b><u>LAB</u></b>	<b><u>PRIORITY</u></b>	<b><u>STUDY DIRECTOR</u></b>	<b><u>CHEMICAL (MFG)</u></b>	<b><u>COMMODITY</u></b>	<b><u>CROP GROUP</u></b>
P12348	-	+	HOMA	FLUTIANIL (LANDIS,OATAGRIO)	SAGE	HERB SUBGROUP (19A)

**Reason for Need:** POWDERY MILDEW - IS NOT BEING CONTROLLED WITH REGISTERED FUNGICIDES

**Use Pattern: (PCR):** MAKE FOLIAR APPLIC OF 0.04 LB AI/A, MINIMUM 7-DAY INTERVAL, MAXIMUM 3 APPLIC/CROP; MFG CONSIDERING REGISTERING A LOWER APPLIC RATE THAN REQUESTED

**E/CS Data Requirements:**

**E/CS Research Comments:**

**IR-4 Residue Trial Plan:**

**Comments:** REQUEST IS FOR USE ON FIELD AND GH-GROWN SAGE; COULD BE COVERED IF DATA IS COLLECTED ON REP CROP BASIL; MFG INDICATES POSSIBLE ASSISTANCE WITH ANALYSIS OF RESIDUE SAMPLES:08/17; MFG CHECKING ON NEED FOR E/CS DATA FIRST:09/17

**NER-EPA Region-FRD**

18-NYP05     Catlin, Nora  
(GH)

**NCR-EPA Region-FRD**

**SOR-EPA Region-FRD**

**WSR-EPA Region-FRD**

**CANADA-EPA Region-FRD**



## 2018 Tentative/Scheduled Studies

### Residue and Efficacy/Crop Safety (E/CS)

*(Order by Crop Group, Commodity, Chemical)*

*Print Date: 2/22/2018*

<b>PR #</b>	<b>LAB</b>	<b>PRIORITY</b>	<b>STUDY DIRECTOR</b>	<b>CHEMICAL (MFG)</b>	<b>COMMODITY</b>	<b>CROP GROUP</b>
P12335	-	+	ARSENOVIC	QUIZALOFOP (DUPONT,GOWAN)	BRASSICA CARINATA	RAPESEED SUBGROUP (20A)

**Reason for Need:** GRASS WEEDS - NO HERBICIDES LABELED IN THE U.S.

**Use Pattern: (PCR):** USE THE ASSURE II PRODUCT; MAKE FOLIAR BROADCAST APPLIC OF 0.047 LB AI/A; CANADIAN LABEL SAYS 64-DAY PHI

**E/CS Data Requirements:**

**E/CS Research Comments:**

**IR-4 Residue Trial Plan:**

**Comments:** CAN BE COVERED WITH RAPESEED CROP SUBGROUP 20A TOLERANCE ESTABLISHED AT 1.5 PPM, IF THE USE PATTERN IS THE SAME AS FOR 20A:08/17; MFG CONFIRMED ONLY NEED E/CS DATA:09/17

**NER-EPA Region-FRD**

**NCR-EPA Region-FRD**

**SOR-EPA Region-FRD**

**WSR-EPA Region-FRD**

**CANADA-EPA Region-FRD**

18-SDP01     Clay, Dr. Sharon  
18-SDP02     Clay, Dr. Sharon



# 2018 Tentative/Scheduled Studies

## Residue and Efficacy/Crop Safety (E/CS)

*(Order by Crop Group, Commodity, Chemical)*

Print Date: 2/22/2018

<b>PR #</b>	<b>LAB</b>	<b>PRIORITY</b>	<b>STUDY DIRECTOR</b>	<b>CHEMICAL (MFG)</b>	<b>COMMODITY</b>	<b>CROP GROUP</b>
12264	18-CAR10	A	DORSCHNER	CYCLANILIPROLE (ISK)	SUNFLOWER	SUNFLOWER SUBGROUP (20B)

**Reason for Need:** TARGET PEST SPECIES ARE LEPIDOPTERAN SPECIES, (HEAD MOTH) AS WELL AS THE NYMPHAL STAGES OF HEMIPTERAN PLANT BUG SPECIES THAT FEED ON DEVELOPING SEEDS IN THE SUNFLOWER SEED HEAD

**Use Pattern: (PCR):** MAKE 3 FOLIAR APPLIC OF 40-80 G AI/HA, 7-DAY INTERVAL, 7-DAY PHI; APPLY IN A MINIMUM 5 GPA BY AIR OR GROUND

**E/CS Data Requirements:** 3 TRIALS; HIGH RATE, LOW RATE, STANDARD, CONTROL (INCLUDE CONFECTIONARY AND BLACK OIL); ALSO EXAMINE PERFORMANCE ON SAFFLOWER

**E/CS Research Comments:**

**IR-4 Residue Trial Plan:** 5-3 7-4 8 (SEED); 1 TRIAL IS FOR PROCESSING (OIL, MEAL); 1 TRIAL IS A DECLNE

**Comments:** THIS IS AN EXPORT COMMODITY, BUT KEY MARKETS NOT IDENTIFIED; ISK MAY CONSIDER FINANCIAL GRANT TO HELP OFFSET COSTS; A RESIDUE STUDY ON SUNFLOWER CAN COVER THE SAFFLOWER NEED IN PR# 11570; ISK SUPPORTS, RESIDUE AND PERFORMANCE DATA NEEDED:07/17

**NER-EPA Region-FRD**

**NCR-EPA Region-FRD**

**SOR-EPA Region-FRD**

**WSR-EPA Region-FRD**

**CANADA-EPA Region-FRD**

18-ND225     Howatt, Kirk  
(reg. 5)(confectionary)  
18-ND226     Howatt, Kirk  
(reg. 5)(decline)(black oil)  
18-ND227     Jenks, Dr. Brian  
(reg. 7)(processing)(black oil)  
18-ND228     Jenks, Dr. Brian  
(reg. 7)(confectionary)  
18-ND229     Jenks, Dr. Brian  
(reg. 7)(black oil)  
18-OH\*270     Horst, Leona  
(seed)(confectionary)  
18-SD333     Clay, Dr. Sharon  
(reg. 5)(black oil)  
18-SD334     Clay, Dr. Sharon  
(reg. 7)(confectionary)

18-CA70     Watkins, S.  
(black oil)  
18-NM258     Hamilton, Cary  
(reg 8)(confectionary)



# 2018 Tentative/Scheduled Studies

## Residue and Efficacy/Crop Safety (E/CS)

*(Order by Crop Group, Commodity, Chemical)*

*Print Date: 2/22/2018*

<b>PR #</b>	<b>LAB</b>	<b>PRIORITY</b>	<b>STUDY DIRECTOR</b>	<b>CHEMICAL (MFG)</b>	<b>COMMODITY</b>	<b>CROP GROUP</b>
P12264	-	A	DORSCHNER	CYCLANILIPROLE (ISK)	SUNFLOWER	SUNFLOWER SUBGROUP (20B)

**Reason for Need:** TARGET PEST SPECIES ARE LEPIDOPTERAN SPECIES, (HEAD MOTH) AS WELL AS THE NYMPHAL STAGES OF HEMIPTERAN PLANT BUG SPECIES THAT FEED ON DEVELOPING SEEDS IN THE SUNFLOWER SEED HEAD

**Use Pattern: (PCR):** MAKE 3 FOLIAR APPLIC OF 40-80 G AI/HA, 7-DAY INTERVAL, 7-DAY PHI; APPLY IN A MINIMUM 5 GPA BY AIR OR GROUND

**E/CS Data Requirements:** 3 TRIALS; HIGH RATE, LOW RATE, STANDARD, CONTROL (INCLUDE CONFECTIONARY AND BLACK OIL); ALSO EXAMINE PERFORMANCE ON SAFFLOWER

**E/CS Research Comments:**

**IR-4 Residue Trial Plan:** 5-3 7-4 8 (SEED); 1 TRIAL IS FOR PROCESSING (OIL, MEAL); 1 TRIAL IS A DECLNE

**Comments:** THIS IS AN EXPORT COMMODITY, BUT KEY MARKETS NOT IDENTIFIED; ISK MAY CONSIDER FINANCIAL GRANT TO HELP OFFSET COSTS; A RESIDUE STUDY ON SUNFLOWER CAN COVER THE SAFFLOWER NEED IN PR# 11570; ISK SUPPORTS, RESIDUE AND PERFORMANCE DATA NEEDED:07/17

**NER-EPA Region-FRD**

**NCR-EPA Region-FRD**

**SOR-EPA Region-FRD**

**WSR-EPA Region-FRD**

**CANADA-EPA Region-FRD**

18-SDP03 Clay, Dr. Sharon  
(likely to be SD)

18-TBD TBD-WSR  
(Kern County potential - safflower)



## 2018 Tentative/Scheduled Studies

### Residue and Efficacy/Crop Safety (E/CS)

*(Order by Crop Group, Commodity, Chemical)*

*Print Date: 2/22/2018*

<b>PR #</b>	<b>LAB</b>	<b>PRIORITY</b>	<b>STUDY DIRECTOR</b>	<b>CHEMICAL (MFG)</b>	<b>COMMODITY</b>	<b>CROP GROUP</b>
P11429	-NONE	A	ARSENOVIC	INDAZIFLAM (BAYER)	ASPARAGUS	STALK AND STEM VEGETABLE SUBGROUP (22A)

**Reason for Need:** ANNUAL WEEDS; DIFFERENT MODE OF ACTION TO AVOID WEED RESISTANCE; CONTROL OF RESISTANT BROADLEAVES

**Use Pattern: (PCR):** 0.065-0.085 LB AI/A OF ALION PRODUCT; ONE PREEMERGENCE (TO THE CROP) APPLIC PER YEAR; 7-DAY PHI; APPLY TO CLEAN SOIL BEFORE ASPARAGUS EMERGES IN THE SPRING

**E/CS Data Requirements:** NEED AT LEAST 2 MORE TRIALS, TESTING 30 D PRE, IN CA AND MI/NJ; ALL SITES REPEAT 2ND YR ON SAME PLOTS:10/17

**E/CS Research Comments:**

**IR-4 Residue Trial Plan:** 2 5-2 10-2 11

**Comments:** MFG APPROVED FOR RESIDUE WORK, BUT REQUIRES ASSESSMENT OF CROP SAFETY WITHIN THE RESIDUE STUDY, ESPECIALLY ON LOW ORGANIC SOILS:09/14; JOINT WITH CANADA (PMC SERVING AS SPONSOR/STUDY DIRECTOR; RESIDUE SITES: 5-3):10/14; MFG NO LONGER CAN SUPPORT THIS PRE-HARVEST USE PATTERN, BUT MIGHT CONSIDER POST-HARVEST USE IF A SAFE USE PATTERN CAN BE FOUND:06/15; RESIDUE STUDY CANCELLED; IR-4 CONTINUING TO SEEK AN ACCEPTABLE POST-HARVEST USE PATTERN:08/15

**NER-EPA Region-FRD**

18-NJP01    Besancon, Thierry

**NCR-EPA Region-FRD**

18-MIP04    Zandstra, Dr. Bernard H.

**SOR-EPA Region-FRD**

**WSR-EPA Region-FRD**

18-CAP09    Hanson, Brad  
(on same plots as 2017 trial in Davis)

**CANADA-EPA Region-FRD**



# 2018 Tentative/Scheduled Studies

## Residue and Efficacy/Crop Safety (E/CS)

(Order by Crop Group, Commodity, Chemical)

Print Date: 2/22/2018

<b>PR #</b>	<b>LAB</b>	<b>PRIORITY</b>	<b>STUDY DIRECTOR</b>	<b>CHEMICAL (MFG)</b>	<b>COMMODITY</b>	<b>CROP GROUP</b>
11830	18-FLR04	A	LENNON	SPINETORAM (DOWAGR)	ASPARAGUS	STALK AND STEM VEGETABLE SUBGROUP (22A)

**Reason for Need:** THRIPS; THEY DAMAGE SPEAR TIPS AND CAUSE THEM TO DRY OUT

**Use Pattern: (PCR):** USE RADIANT SC PRODUCT; MAKE 6 FOLIAR APPLIC TO SPEARS OF 6-10 OZ PROD/A AT 4-DAY INTERVALS IN MINIMUM 10 GPA BY AIR OR 20 GPA BY GROUND, IN-SEASON DURING SPEAR HARVESTING; 1-DAY PHI; USE OF ADJUVANTS OR TANK-MIXING WITH EC PRODUCTS COULD CAUSE UNACCEPTABLE APPEARANCE; PCR USE PATTERN UPDATE FROM REQUESTOR 03/28/17: MAX 4 APPLIC BETWEEN JAN. 1 AND JUNE 1 IS ACCEPTABLE, AND A 1-DAY PHI IS PREFERRED

**E/CS Data Requirements:**

**E/CS Research Comments:**

**IR-4 Residue Trial Plan:** 2 5-2 10-3 11-2 (1 DECLINE TRIAL)

**Comments:** LIKELY EXPORT MARKETS WERE NOT DEFINED; THERE IS A SPINETORAM TOLERANCE ON ASPARAGUS, BUT DOES NOT APPEAR TO BE LABELED; LIKELY REQUIRES MOR STUDY AND PERFORMANCE DATA:10/15; IS AN "A" PRIORITY IN CANADA FOR 2017, SO POTENTIAL JOINT PROJECT:06/16; EPA CAUTION:08/16; EPA GREEN:08/17

**NER-EPA Region-FRD**

18-NJ240      Freiburger, Tom

**NCR-EPA Region-FRD**

18-MI188      Zandstra, Dr. Bernard H.  
18-MI189      Zandstra, Dr. Bernard H.

**SOR-EPA Region-FRD**

**WSR-EPA Region-FRD**

18-CA39      Skiles, Keri  
18-CA40      Kyser, Guy  
(decline)  
18-CA\*41      Benzen, Ms. Sharon D.  
18-WA351      Peng, Wilson  
18-WA352      Peng, Wilson

**CANADA-EPA Region-FRD**



# 2018 Tentative/Scheduled Studies

## Residue and Efficacy/Crop Safety (E/CS)

(Order by Crop Group, Commodity, Chemical)

Print Date: 2/22/2018

<u>PR #</u>	<u>LAB</u>	<u>PRIORITY</u>	<u>STUDY DIRECTOR</u>	<u>CHEMICAL (MFG)</u>	<u>COMMODITY</u>	<u>CROP GROUP</u>
P10184	-NONE	A	BARNEY	RIMSULFURON (DUPONT)	OLIVE	TROPICAL AND SUBTROPICAL, SMALL FRUIT, EDIBLE PEEL SUBGROUP (23A)

**Reason for Need:** ANNUAL BROADLEAVES & GRASSES, INCLUDING HAIRY FLEABANE, MARESTAIL, RYEGRASS, PANICLE WILLOWHERB & PUNCTUREVINE

**Use Pattern: (PCR):** 1 OZ AI/A; GROUND APPLIC TO THE SOIL; 1 OR 2 APPLIC; 30-DAY RE-TREATMENT INTERVAL; 14-DAY PHI; APPLY PRIOR TO WEED EMERGENCE; NO MORE THAN 4 OZ OF PRODUCT APPLIED PER YEAR BROADCAST, OR 8 OZ OF PRODUCT IF APPLIED IN A BAND OF LESS THAN 50% OF THE TOTAL ORCHARD

**E/CS Data Requirements:** 3-4 TRIALS; MFG CONFIRMED THAT CROP SAFETY DATA SHOULD BE DOCUMENTED DURING CONDUCT OF RESIDUE TRIALS, AND THAT EFFICACY DATA B

**E/CS Research Comments:** IN 2017 PERFORMANCE PROTOCOL: TESTING RIMSULFURON (MATRIX) AT 0.0625 AND 0.125 LB AI/A, IN 10 OR MORE GPA, INCLUDING ADJUVANT AT A LABELED RATE, COMPARED WITH A STANDARD; MAKE 2 SOIL BANDED APPLIC TO THE ORCHARD FLOOR ON EACH SIDE OF THE TREE ROW, 30-DAY INTERVAL, WITH LAST APPLIC 14 DAYS BEFORE HARVEST; COLLECT DATA ON CROP INJURY, WEED CONTROL AND CROP YIELD

**IR-4 Residue Trial Plan:** 3/6 10-4, 1 PROCESSING TRIAL (OIL) (NEED MIN. 5 TRIALS FOR U.S., 8 FOR CODEX)

**Comments:** MFG REQUESTS THAT WEED CONTROL AND CROP TOLERANCE RATINGS BE TAKEN DURING RESIDUE STUDY; ANY WEEDS TO BE LISTED ON LABEL MUST BE SUPPORTED BY EFFICACY DATA:07/10; MFG MAY HAVE SOME CS DATA:06/14; MFG TO CONFIRM PROJECT STATUS AFTER ASSESSING CA DATA:06/15; AT 2015 FUW, MFG CONFIRMED THIS IS RESEARCHABLE, RESIDUE AND E/CS DATA:09/15

[NER-EPA Region-FRD](#)

[NCR-EPA Region-FRD](#)

[SOR-EPA Region-FRD](#)

[WSR-EPA Region-FRD](#)

[CANADA-EPA Region-FRD](#)

18-CAP10 Hanson, Brad





## 2018 Tentative/Scheduled Studies

### Residue and Efficacy/Crop Safety (E/CS)

*(Order by Crop Group, Commodity, Chemical)*

*Print Date: 2/22/2018*

<u>PR #</u>	<u>LAB</u>	<u>PRIORITY</u>	<u>STUDY DIRECTOR</u>	<u>CHEMICAL (MFG)</u>	<u>COMMODITY</u>	<u>CROP GROUP</u>
12404	18-TBD	A	SAMOIL	SULFUR DIOXIDE (TEDMRK)	FIG	TROPICAL AND SUBTROPICAL, MEDIUM TO LARGE FRUIT, EDIBLE PEEL SUBGROUP (23B)

**Reason for Need:** ASPERGILLUS MOLD, BOTRYTIS FRUIT ROT, SOFT ROT, SURFACE MOLD

**Use Pattern: (PCR):** FROM ORIGINAL REQUEST 10114: 25-100 PPM-HOUR; GAS APPLIC AFTER HARVEST (25-100 PPM-HOUR), SULFUR DIOXIDE PAD (0.2-25 PPM); 1 APPLIC

**E/CS Data Requirements:**

**E/CS Research Comments:**

**IR-4 Residue Trial Plan:** 10-4

**Comments:** THIS IS A NEW PR# FOR A REPEAT STUDY, AFTER THE PETITION FROM THE ORIGINAL STUDY PR#10114 (2009) HAD TO BE WITHDRAWN FROM EPA:10/17

**NER-EPA Region-FRD**

**NCR-EPA Region-FRD**

**SOR-EPA Region-FRD**

**WSR-EPA Region-FRD**

**CANADA-EPA Region-FRD**

- 18-CA98 Ennes, D. (Kearney)  
(special equipment needed)
- 18-CA99 Ennes, D. (Kearney)  
(special equipment needed)
- 18-CA100 Ennes, D. (Kearney)  
(special equipment needed)
- 18-CA101 Ennes, D. (Kearney)  
(special equipment needed)



## 2018 Tentative/Scheduled Studies

### Residue and Efficacy/Crop Safety (E/CS)

*(Order by Crop Group, Commodity, Chemical)*

*Print Date: 2/22/2018*

<u>PR #</u>	<u>LAB</u>	<u>PRIORITY</u>	<u>STUDY DIRECTOR</u>	<u>CHEMICAL (MFG)</u>	<u>COMMODITY</u>	<u>CROP GROUP</u>
10242	18-TBD	A	ARSENOVIC	GLUFOSINATE (BAYER,UPI)	GUAVA	TROPICAL AND SUBTROPICAL, MEDIUM TO LARGE FRUIT, EDIBLE PEEL SUBGROUP (23B)

**Reason for Need:** PARTHENIUM HYSTEROPHORUS

**Use Pattern: (PCR):** 10.5 OZ/A X BAND WIDTH; BANDED APPLIC; 6 APPLIC; 30-60 RE-TREATMENT INTERVALS; 0-DAY PHI; CALCULATE BANDWIDTH TO HERBICIDE, THEN MIX SUFFICIENT RELY WITH 30-40 GAL/WATER/TREATED/A AND MAY BE MIXED; DO NOT APPLY MORE THAN 345 FL OZ RELY/A/YEAR; MFG REQUESTS SAME USE PATTERN AS ON FIG:05/17

**E/CS Data Requirements:**

**E/CS Research Comments:**

**IR-4 Residue Trial Plan:** 13-4

**Comments:** ORIGINAL REQUEST REC'D 8/25/2008; MFG NO - THERE ARE RISK CUP & GROUND WATER CONCERNS:08/08; MFG WILL REVISIT AFTER RE-REG REVIEW IS COMPLETED BY EPA:05/16; MFG SUPPORTS (RESIDUE + E/CS DATA); ALONG WITH FIG (PR# 11547) IS THE OTHER REP CROP FOR NEW SUBGROUP 23B:08/16; EPA CAUTION:09/16; EPA CAUTION:08/17

**NER-EPA Region-FRD**

**NCR-EPA Region-FRD**

**SOR-EPA Region-FRD**

**WSR-EPA Region-FRD**

**CANADA-EPA Region-FRD**

18-FL113	Tannenbaum, Rebecca	18-CA27	Leach, Nathan
18-PR308	Robles Vazquez, W.	18-HI149	Kam, James



## 2018 Tentative/Scheduled Studies Residue and Efficacy/Crop Safety (E/CS)

(Order by Crop Group, Commodity, Chemical)

Print Date: 2/22/2018

<u>PR #</u>	<u>LAB</u>	<u>PRIORITY</u>	<u>STUDY DIRECTOR</u>	<u>CHEMICAL (MFG)</u>	<u>COMMODITY</u>	<u>CROP GROUP</u>
P10242	-	A	ARSENOVIC	GLUFOSINATE (BAYER,UPI)	GUAVA	TROPICAL AND SUBTROPICAL, MEDIUM TO LARGE FRUIT, EDIBLE PEEL SUBGROUP (23B)

**Reason for Need:** PARTHENIUM HYSTEROPHORUS

**Use Pattern: (PCR):** 10.5 OZ/A X BAND WIDTH; BANDED APPLIC; 6 APPLIC; 30-60 RE-TREATMENT INTERVALS; 0-DAY PHI; CALCULATE BANDWIDTH TO HERBICIDE, THEN MIX SUFFICIENT RELY WITH 30-40 GAL/WATER/TREATED/A AND MAY BE MIXED; DO NOT APPLY MORE THAN 345 FL OZ RELY/A/YEAR; MFG REQUESTS SAME USE PATTERN AS ON FIG:05/17

**E/CS Data Requirements:**

**E/CS Research Comments:**

**IR-4 Residue Trial Plan:** 13-4

**Comments:** ORIGINAL REQUEST REC'D 8/25/2008; MFG NO - THERE ARE RISK CUP & GROUND WATER CONCERNS:08/08; MFG WILL REVISIT AFTER RE-REG REVIEW IS COMPLETED BY EPA:05/16; MFG SUPPORTS (RESIDUE + E/CS DATA); ALONG WITH FIG (PR# 11547) IS THE OTHER REP CROP FOR NEW SUBGROUP 23B:08/16; EPA CAUTION:09/16; EPA CAUTION:08/17

**NER-EPA Region-FRD**

**NCR-EPA Region-FRD**

**SOR-EPA Region-FRD**

**WSR-EPA Region-FRD**

**CANADA-EPA Region-FRD**

18-FLP06 Crane, Dr. Jonathan H.



## 2018 Tentative/Scheduled Studies

### Residue and Efficacy/Crop Safety (E/CS)

*(Order by Crop Group, Commodity, Chemical)*

*Print Date: 2/22/2018*

<u>PR #</u>	<u>LAB</u>	<u>PRIORITY</u>	<u>STUDY DIRECTOR</u>	<u>CHEMICAL (MFG)</u>	<u>COMMODITY</u>	<u>CROP GROUP</u>
P8540	-NONE	+	SAMOIL	BIFENTHRIN (ADAMA,AMVAC,FMC)	LYCHEE	TROPICAL AND SUBTROPICAL, SMALL FRUIT, INEDIBLE PEEL SUBGROUP (24A)

**Reason for Need:** MYLLOCERUS WEEVIL

**Use Pattern: (PCR):** SEE PRODUCT LABEL DIRECTIONS FOR CITRUS, FOR CAPTURE SOIL PRODUCT AND FURY FOLIAR PRODUCT; FMC REQUESTS USE OF WSB PRODUCT:06/16

**E/CS Data Requirements:** NEED 1 GOOD TRIAL, WITH FOLIAR TRT, SOIL TRT (BARRIER TRT), FOLIAR+SOIL TRT, UNTREATED:10/17

**E/CS Research Comments:**

**IR-4 Residue Trial Plan:**

**Comments:** MFG REQUIRES EFFICACY DATA PRIOR TO RESIDUE STUDY:07/03; MFG CONSIDERING DOING E/CS RESEARCH:05/13; EPA CAUTION:08/15; EPA CAUTION:09/16

**NER-EPA Region-FRD**

**NCR-EPA Region-FRD**

**SOR-EPA Region-FRD**

**WSR-EPA Region-FRD**

**CANADA-EPA Region-FRD**

18-FLP11 Crane, Dr. Jonathan H.  
(FL-Crane?)



# 2018 Tentative/Scheduled Studies

## Residue and Efficacy/Crop Safety (E/CS)

(Order by Crop Group, Commodity, Chemical)

Print Date: 2/22/2018

<u>PR #</u>	<u>LAB</u>	<u>PRIORITY</u>	<u>STUDY DIRECTOR</u>	<u>CHEMICAL (MFG)</u>	<u>COMMODITY</u>	<u>CROP GROUP</u>
10239	18-TBD	A	ARSENOVIC	GLUFOSINATE (BAYER,UPI)	LYCHEE	TROPICAL AND SUBTROPICAL, SMALL FRUIT, INEDIBLE PEEL SUBGROUP (24A)

**Reason for Need:** PARTHENIUM HYSTEROPHORUS

**Use Pattern: (PCR):** 10.5 OZ/A X BAND WIDTH; BANDED APPLIC; 6 APPLIC; 30-60 RE-TREATMENT INTERVALS; 0-DAY PHI; CALCULATE BANDWIDTH TO HERBICIDE, THEN MIX SUFFICIENT RELY WITH 30-40 GAL/WATER/TREATED/A AND MAY BE MIXED; DO NOT APPLY MORE THAN 345 FL OZ RELY/A/YEAR; MFG SUGGESTS COMMON USE PATTERN FOR ALL TROPICAL FRUITS:05/17

**E/CS Data Requirements:**

**E/CS Research Comments:**

**IR-4 Residue Trial Plan:** 13-4

**Comments:** ORIGINAL REQUEST REC'D 8/25/2008; MFG NO - THERE ARE RISK CUP & GROUND WATER CONCERNS:08/08; MFG WILL REVISIT AFTER RE-REG REVIEW IS COMPLETED BY EPA:05/16; MFG SUPPORTS (RESIDUE + E/CS DATA); IS REP CROP FOR NEW SUBGROUP 24A:08/16; EPA CAUTION:09/16; EPA CAUTION:08/17

**NER-EPA Region-FRD**

**NCR-EPA Region-FRD**

**SOR-EPA Region-FRD**

**WSR-EPA Region-FRD**

**CANADA-EPA Region-FRD**

18-FL111	Tannenbaum, Rebecca	18-HI148	Kam, James
18-FL112	Tannenbaum, Rebecca		
18-PR307	Robles Vazquez, W.		



## 2018 Tentative/Scheduled Studies

### Residue and Efficacy/Crop Safety (E/CS)

*(Order by Crop Group, Commodity, Chemical)*

*Print Date: 2/22/2018*

<u>PR #</u>	<u>LAB</u>	<u>PRIORITY</u>	<u>STUDY DIRECTOR</u>	<u>CHEMICAL (MFG)</u>	<u>COMMODITY</u>	<u>CROP GROUP</u>
P10239	-	A	ARSENOVIC	GLUFOSINATE (BAYER,UPI)	LYCHEE	TROPICAL AND SUBTROPICAL, SMALL FRUIT, INEDIBLE PEEL SUBGROUP (24A)

**Reason for Need:** PARTHENIUM HYSTEROPHORUS

**Use Pattern: (PCR):** 10.5 OZ/A X BAND WIDTH; BANDED APPLIC; 6 APPLIC; 30-60 RE-TREATMENT INTERVALS; 0-DAY PHI; CALCULATE BANDWIDTH TO HERBICIDE, THEN MIX SUFFICIENT RELY WITH 30-40 GAL/WATER/TREATED/A AND MAY BE MIXED; DO NOT APPLY MORE THAN 345 FL OZ RELY/A/YEAR; MFG SUGGESTS COMMON USE PATTERN FOR ALL TROPICAL FRUITS:05/17

**E/CS Data Requirements:**

**E/CS Research Comments:**

**IR-4 Residue Trial Plan:** 13-4

**Comments:** ORIGINAL REQUEST REC'D 8/25/2008; MFG NO - THERE ARE RISK CUP & GROUND WATER CONCERNS:08/08; MFG WILL REVISIT AFTER RE-REG REVIEW IS COMPLETED BY EPA:05/16; MFG SUPPORTS (RESIDUE + E/CS DATA); IS REP CROP FOR NEW SUBGROUP 24A:08/16; EPA CAUTION:09/16; EPA CAUTION:08/17

**NER-EPA Region-FRD**

**NCR-EPA Region-FRD**

**SOR-EPA Region-FRD**

**WSR-EPA Region-FRD**

**CANADA-EPA Region-FRD**

18-FLP07 Crane, Dr. Jonathan H.



# 2018 Tentative/Scheduled Studies

## Residue and Efficacy/Crop Safety (E/CS)

*(Order by Crop Group, Commodity, Chemical)*

*Print Date: 2/22/2018*

<u>PR #</u>	<u>LAB</u>	<u>PRIORITY</u>	<u>STUDY DIRECTOR</u>	<u>CHEMICAL (MFG)</u>	<u>COMMODITY</u>	<u>CROP GROUP</u>
P10240	-NONE	A	ARSENOVIC	GLUFOSINATE (BAYER,UPI)	AVOCADO	TROPICAL AND SUBTROPICAL, MEDIUM TO LARGE FRUIT, SMOOTH, INEDIBLE PEEL SUBGROUP (24B)

**Reason for Need:** PARTHENIUM HYSTEROPHORUS

**Use Pattern: (PCR):** 10.5 OZ/A X BAND WIDTH; BANDED APPLIC; 6 APPLIC; 30-60 RE-TREATMENT INTERVALS; 0-DAY PHI; CALCULATE BANDWIDTH TO HERBICIDE, THEN MIX SUFFICIENT RELY WITH 30-40 GAL/WATER/TREATED/A AND MAY BE MIXED; DO NOT APPLY MORE THAN 345 FL OZ RELY/A/YEAR

**E/CS Data Requirements:** 3-5 TRIALS; IN 2018 NEED 2 TRIALS WITH NEW USE PATTERN LANGUAGE; LABEL LIKELY TO REQUIRE LANGUAGE SPECIFYING DROPLET SIZE:10/17

**E/CS Research Comments:** IN THE 2017 PROTOCOL, TESTING 2 RATES OF GLUFOSINATE (RELY 280, 2.34 LB AI/GAL) ALONE (1.5 AND 3.0 LB AI/A) AND THE SAME RATES TANK-MIXED WITH A LABELED RATE OF GLYPHOSATE; APPLIC ARE BANDED TO THE ORCHARD FLOOR IN >15 GPA, 2 APPLIC, 28 DAYS APART, WITH FIRST APPLIC WHEN WEEDS ARE <6 IN TALL; ADJUVANT IS INCLUDED WITH GLUFOSINATE ALONE; COLLECTING CROP INJURY AND WEED CONTROL DATA

**IR-4 Residue Trial Plan:** 13-2 10-3, 1 DECLINE TRIAL

**Comments:** ORIGINAL REQUEST REC'D 8/25/2008; MFG NO - THERE ARE RISK CUP & GROUND WATER CONCERNS:08/08; MFG WILL REVISIT AFTER RE-REG REVIEW IS COMPLETED BY EPA:05/16; MFG HAS COMPLETED TRIALS FOR IMPORT TOLERANCE THAT MIGHT HELP REDUCE # OF RESIDUE TRIALS NEEDED; THERE ARE ESTABLISHED EPA TOLERANCES FOR BANANA (0.3 PPM) AND BANANA PULP (0.2 PPM); NEED TO HARMONIZE USE PATTERNS SO THAT BANANA (PR# 12050) AND AVOCADO, AS REP CROPS FOR NEW SUBGROUP 24B, CAN COVER PAPAYA (PR# 09887):08/16; MFG SUPPORTS (RESIDUE + E/CS DATA); EPA CAUTION:09/16; MIR IS STORING SAMPLES UNTIL A CONTRACT LAB IS ASSIGNED: 9/17

**NER-EPA Region-FRD**

**NCR-EPA Region-FRD**

**SOR-EPA Region-FRD**

**WSR-EPA Region-FRD**

**CANADA-EPA Region-FRD**

18-CAP11 Rios, Sonia  
18-CAP12 Rios, Sonia



## 2018 Tentative/Scheduled Studies Residue and Efficacy/Crop Safety (E/CS)

(Order by Crop Group, Commodity, Chemical)

Print Date: 2/22/2018

<b>PR #</b>	<b>LAB</b>	<b>PRIORITY</b>	<b>STUDY DIRECTOR</b>	<b>CHEMICAL (MFG)</b>	<b>COMMODITY</b>	<b>CROP GROUP</b>
07337	18-TIR01	A	BARNEY	TEBUCONAZOLE (ADAMA,BAYER)	AVOCADO	TROPICAL AND SUBTROPICAL, MEDIUM TO LARGE FRUIT, SMOOTH, INEDIBLE PEEL SUBGROUP (24B)

**Reason for Need:** FRUIT ROT, LEAF & FRUIT SPOTS

**Use Pattern: (PCR):** NO USE PATTERN PROVIDED CONTACTED WSR:04/10

**E/CS Data Requirements:**

**E/CS Research Comments:**

**IR-4 Residue Trial Plan:** 3-2 10-3; 1 TRIAL IS A DECLINE

**Comments:** NEED ANTICIPATED RESIDUE DATA:08/00; DROPPED DUE TO 3 YEAR NON-NOMINATION:08/12; ADAMA SUPPORTS, AT FUW; NEED SUBSTITUTES FOR COPPER SPRAYS TO CONTROL SCAB AND ANTHRACNOSE:09/17; EPA CAUTION:11/17

**NER-EPA Region-FRD**

**NCR-EPA Region-FRD**

**SOR-EPA Region-FRD**

**WSR-EPA Region-FRD**

**CANADA-EPA Region-FRD**

18-FL105	Tannenbaum, Rebecca	18-CA11	Skiles, Keri
18-FL106	Tannenbaum, Rebecca	18-CA12	Leach, Nathan
18-PR306	Robles Vazquez, W.	(decline)	
		18-CA13	Leach, Nathan





## 2018 Tentative/Scheduled Studies

### Residue and Efficacy/Crop Safety (E/CS)

*(Order by Crop Group, Commodity, Chemical)*

*Print Date: 2/22/2018*

<u>PR #</u>	<u>LAB</u>	<u>PRIORITY</u>	<u>STUDY DIRECTOR</u>	<u>CHEMICAL (MFG)</u>	<u>COMMODITY</u>	<u>CROP GROUP</u>
11160	16-TIR01	A	JOLLY	TEBUCONAZOLE (ADAMA,BAYER)	AVOCADO	TROPICAL AND SUBTROPICAL, MEDIUM TO LARGE FRUIT, SMOOTH, INEDIBLE PEEL SUBGROUP (24B)

**Reason for Need:** NEED CONTROL OF DISEASE CAUSED BY R. LAURICOLA

**Use Pattern: (PCR):** 20-40 ML/INCH TRUNK DIAMETER; INJECTION OF INFUSION INTO SAPWOOD; 1 APPLIC/YEAR; 1 YEAR RE-TREATMENT INTERVAL; 1-DAY PHI; DO NOT APPLY MORE THAN 60 ML/INCH TRUNK DIAMETER

**E/CS Data Requirements:**

**E/CS Research Comments:**

**IR-4 Residue Trial Plan:** 3-2 10-3 (1 DECLINE)

**Comments:** BAYER HAS NO SOLO TEBUCONAZOLE PRODUCT; DISCUSS WITH MANA FOR FOLICUR:04/13; AT FUW BOTH MFG INDICATED WILL NOT SUPPORT:09/13; ADAMA WILL SUPPORT THIS REQUEST:07/15; EPA CAUTION:08/15

**NER-EPA Region-FRD**

**NCR-EPA Region-FRD**

**SOR-EPA Region-FRD**

**WSR-EPA Region-FRD**

**CANADA-EPA Region-FRD**

18-CA1      Leach, Nathan  
(2018 \$)



# 2018 Tentative/Scheduled Studies

## Residue and Efficacy/Crop Safety (E/CS)

(Order by Crop Group, Commodity, Chemical)

Print Date: 2/22/2018

<u>PR #</u>	<u>LAB</u>	<u>PRIORITY</u>	<u>STUDY DIRECTOR</u>	<u>CHEMICAL (MFG)</u>	<u>COMMODITY</u>	<u>CROP GROUP</u>
12050	18-TBD	A	ARSENOVIC	GLUFOSINATE (BAYER,UPI)	BANANA	TROPICAL AND SUBTROPICAL, MEDIUM TO LARGE FRUIT, SMOOTH, INEDIBLE PEEL SUBGROUP (24B)

**Reason for Need:** VARIOUS BROADLEAF AND GRASSY WEEDS

**Use Pattern: (PCR):** USE THE RELY 280 PRODUCT; MAKE 3 DIRECTED APPLIC/YEAR OF 1.5 LB A/A, 30-DAY INTERVAL, 0-DAY PHI; APPLY OVER ROW MIDDLES OR SPOT SPRAY; MFG SUGGESTS COMMON USE PATTERN FOR ALL TROPICAL FRUITS:05/17

**E/CS Data Requirements:**

**E/CS Research Comments:**

**IR-4 Residue Trial Plan:** 3-2 13-6 (2-YR STUDY); 2 "RED A" TRIALS NEEDED IN 2019

**Comments:** THERE ARE ESTABLISHED EPA TOLERANCES FOR BANANA (0.3 PPM) AND BANANA PULP (0.2 PPM); NEED TO HARMONIZE USE PATTERNS SO THAT AVOCADO (PR# 10240) AND BANANA, AS REP CROPS FOR NEW SUBGROUP 24B, WILL COVER PAPAYA (PR# 09887):08/16; EPA CAUTION; MFG SUPPORTS, BUT NEED E/CS DATA BEFORE RESIDUE STUDY:09/16; MFG SUPPORTS, WITH E/CS DATA GENERATED DURING RESIDUE STUDY:05/17; EPA CAUTION:11/17

**NER-EPA Region-FRD**

**NCR-EPA Region-FRD**

**SOR-EPA Region-FRD**

**WSR-EPA Region-FRD**

**CANADA-EPA Region-FRD**

18-FL122	Tannenbaum, Rebecca	18-HI153	Coughlin, Julie
18-FL123	Tannenbaum, Rebecca	18-HI154	Coughlin, Julie
(bagged)		18-HI155	Coughlin, Julie
		18-HI156	Coughlin, Julie
		(bagged)	



## 2018 Tentative/Scheduled Studies Residue and Efficacy/Crop Safety (E/CS)

(Order by Crop Group, Commodity, Chemical)

Print Date: 2/22/2018

<u>PR #</u>	<u>LAB</u>	<u>PRIORITY</u>	<u>STUDY DIRECTOR</u>	<u>CHEMICAL (MFG)</u>	<u>COMMODITY</u>	<u>CROP GROUP</u>
P12050	-	A	ARSENOVIC	GLUFOSINATE (BAYER,UPI)	BANANA	TROPICAL AND SUBTROPICAL, MEDIUM TO LARGE FRUIT, SMOOTH, INEDIBLE PEEL SUBGROUP (24B)

**Reason for Need:** VARIOUS BROADLEAF AND GRASSY WEEDS

**Use Pattern: (PCR):** USE THE RELY 280 PRODUCT; MAKE 3 DIRECTED APPLIC/YEAR OF 1.5 LB A/A, 30-DAY INTERVAL, 0-DAY PHI; APPLY OVER ROW MIDDLES OR SPOT SPRAY; MFG SUGGESTS COMMON USE PATTERN FOR ALL TROPICAL FRUITS:05/17

**E/CS Data Requirements:**

**E/CS Research Comments:**

**IR-4 Residue Trial Plan:** 3-2 13-6 (2-YR STUDY); 2 "RED A" TRIALS NEEDED IN 2019

**Comments:** THERE ARE ESTABLISHED EPA TOLERANCES FOR BANANA (0.3 PPM) AND BANANA PULP (0.2 PPM); NEED TO HARMONIZE USE PATTERNS SO THAT AVOCADO (PR# 10240) AND BANANA, AS REP CROPS FOR NEW SUBGROUP 24B, WILL COVER PAPAYA (PR# 09887):08/16; EPA CAUTION; MFG SUPPORTS, BUT NEED E/CS DATA BEFORE RESIDUE STUDY:09/16; MFG SUPPORTS, WITH E/CS DATA GENERATED DURING RESIDUE STUDY:05/17; EPA CAUTION:11/17

**NER-EPA Region-FRD**

**NCR-EPA Region-FRD**

**SOR-EPA Region-FRD**

**WSR-EPA Region-FRD**

**CANADA-EPA Region-FRD**

18-FLP08 Crane, Dr. Jonathan H.



# 2018 Tentative/Scheduled Studies

## Residue and Efficacy/Crop Safety (E/CS)

(Order by Crop Group, Commodity, Chemical)

Print Date: 2/22/2018

<u>PR #</u>	<u>LAB</u>	<u>PRIORITY</u>	<u>STUDY DIRECTOR</u>	<u>CHEMICAL (MFG)</u>	<u>COMMODITY</u>	<u>CROP GROUP</u>
11572	17-TIR03	A	JOLLY	DIFENOCONAZOLE + AZOXYSTROBIN (SYNGEN)	MANGO	TROPICAL AND SUBTROPICAL, MEDIUM TO LARGE FRUIT, SMOOTH, INEDIBLE PEEL SUBGROUP (24B)

**Reason for Need:** COLLETOTRICHIM SP/ANTHRACNOSE; BLOSSOM BLIGHT & FRUIT ROT

**Use Pattern: (PCR):** 16-20 OZ/A OF INSPIRE PRODUCT; 15 FOLIAR APPLIC, STARTING AT BEGINNING OF FLOWERING THROUGH HARVEST; 7-DAY RE-TREATMENT INTERVAL; 0-DAY PHI

**E/CS Data Requirements:**

**E/CS Research Comments:** IN 2017 PERF. PROTOCOL, TESTING QUADRIS TOP (AZOXY + DIFEN) VS SWITCH (CYPRODINIL + FLUDIOXONIL); MAKE 4 FOLIAR DIRECTED, AIRBLAST OR HANDGUN APPLIC OF EACH TREATMENT, 7-DAY INTERVALS, IN 80-200 GPA, STARTING WHEN CONDITIONS ARE FAVORABLE FOR ANTHRACNOSE DEVELOPMENT; INCLUDE AN NIS ADJUVANT AT A LABELED RATE WITH EACH APPLIC; COLLECTING CROP SAFETY AND PEST CONTROL (ON FRUIT AND FOLIAGE) DATA

**IR-4 Residue Trial Plan:** 13-4 (DOING 2 TRIALS IN 2017; 2 "RED A" TRIALS IN 2018, WITH 1 DECLINE)

**Comments:** TOLERANCE IS ESTABLISHED FOR IMPORTED MANGO; SOME DATA IS AVAILABLE FROM BRAZIL:08/11; USE OF DIFENOCONAZOLE WAS ORIGINALLY REQUESTED ON MANGO, BUT RESIDUE STUDY WAS RUN ON PAPAYA (PR# 10802) WHICH, BY EPA DEFINITION, COVERED MANGO; NOW MANGO IS COVERED BY REP CROPS AVOCADO, PLUS POMEGRANATE OR BANANA IN NEW CROP SUBGROUP 24B, AND MANGO IS NOT COVERED BY PAPAYA DATA; WITH TROPICAL FRUIT CROP GROUPS NOW CODIFIED, USE ON MANGO NEEDS TO BE SUPPORTED WITH RESIDUE DATA ON MANGO, OR COVERED BY DATA FOR THE SUBGROUP 24B REP CROPS:05/16; RESIDUE DATA ARE BEING GENERATED USING THE DUAL AI PRODUCT DIFEN + AZOXYSTROBIN, SO REQUESTED CHEMICAL WAS CHANGED FROM DIFENOCONAZOLE-ONLY TO THIS DUAL AI PRODUCT:11/16

**NER-EPA Region-FRD**

**NCR-EPA Region-FRD**

**SOR-EPA Region-FRD**

**WSR-EPA Region-FRD**

**CANADA-EPA Region-FRD**

18-FL116 Tannenbaum, Rebecca  
 18-FL117 Tannenbaum, Rebecca  
 (decline)



## 2018 Tentative/Scheduled Studies Residue and Efficacy/Crop Safety (E/CS)

(Order by Crop Group, Commodity, Chemical)

Print Date: 2/22/2018

<u>PR #</u>	<u>LAB</u>	<u>PRIORITY</u>	<u>STUDY DIRECTOR</u>	<u>CHEMICAL (MFG)</u>	<u>COMMODITY</u>	<u>CROP GROUP</u>
P11572	-	A	HOMA	DIFENOCONAZOLE + AZOXYSTROBIN (SYNGEN)	MANGO	TROPICAL AND SUBTROPICAL, MEDIUM TO LARGE FRUIT, SMOOTH, INEDIBLE PEEL SUBGROUP (24B)

**Reason for Need:** COLLETOTRICHIM SP/ANTHRACNOSE; BLOSSOM BLIGHT & FRUIT ROT

**Use Pattern: (PCR):** 16-20 OZ/A OF INSPIRE PRODUCT; 15 FOLIAR APPLIC, STARTING AT BEGINNING OF FLOWERING THROUGH HARVEST; 7-DAY RE-TREATMENT INTERVAL; 0-DAY PHI

**E/CS Data Requirements:**

**E/CS Research Comments:** IN 2017 PERF. PROTOCOL, TESTING QUADRIS TOP (AZOXY + DIFEN) VS SWITCH (CYPRODINIL + FLUDIOXONIL); MAKE 4 FOLIAR DIRECTED, AIRBLAST OR HANDGUN APPLIC OF EACH TREATMENT, 7-DAY INTERVALS, IN 80-200 GPA, STARTING WHEN CONDITIONS ARE FAVORABLE FOR ANTHRACNOSE DEVELOPMENT; INCLUDE AN NIS ADJUVANT AT A LABELED RATE WITH EACH APPLIC; COLLECTING CROP SAFETY AND PEST CONTROL (ON FRUIT AND FOLIAGE) DATA

**IR-4 Residue Trial Plan:** 13-4 (DOING 2 TRIALS IN 2017; 2 "RED A" TRIALS IN 2018, WITH 1 DECLINE)

**Comments:** TOLERANCE IS ESTABLISHED FOR IMPORTED MANGO; SOME DATA IS AVAILABLE FROM BRAZIL:08/11; USE OF DIFENOCONAZOLE WAS ORIGINALLY REQUESTED ON MANGO, BUT RESIDUE STUDY WAS RUN ON PAPAYA (PR# 10802) WHICH, BY EPA DEFINITION, COVERED MANGO; NOW MANGO IS COVERED BY REP CROPS AVOCADO, PLUS POMEGRANATE OR BANANA IN NEW CROP SUBGROUP 24B, AND MANGO IS NOT COVERED BY PAPAYA DATA; WITH TROPICAL FRUIT CROP GROUPS NOW CODIFIED, USE ON MANGO NEEDS TO BE SUPPORTED WITH RESIDUE DATA ON MANGO, OR COVERED BY DATA FOR THE SUBGROUP 24B REP CROPS:05/16; RESIDUE DATA ARE BEING GENERATED USING THE DUAL AI PRODUCT DIFEN + AZOXYSTROBIN, SO REQUESTED CHEMICAL WAS CHANGED FROM DIFENOCONAZOLE-ONLY TO THIS DUAL AI PRODUCT:11/16

**NER-EPA Region-FRD**

**NCR-EPA Region-FRD**

**SOR-EPA Region-FRD**

**WSR-EPA Region-FRD**

**CANADA-EPA Region-FRD**

18-FLP09 Crane, Dr. Jonathan H.



## 2018 Tentative/Scheduled Studies Residue and Efficacy/Crop Safety (E/CS)

*(Order by Crop Group, Commodity, Chemical)*

*Print Date: 2/22/2018*

<u>PR #</u>	<u>LAB</u>	<u>PRIORITY</u>	<u>STUDY DIRECTOR</u>	<u>CHEMICAL (MFG)</u>	<u>COMMODITY</u>	<u>CROP GROUP</u>
08274	16-CAR18	A	BARNEY	FLUAZINAM (ISK,SYNGEN)	PAPAYA	TROPICAL AND SUBTROPICAL, MEDIUM TO LARGE FRUIT, SMOOTH, INEDIBLE PEEL SUBGROUP (24B)

**Reason for Need:** ANTHRACNOSE

**Use Pattern: (PCR):** GROUND; 2 APPLIC; 14-DAY PHI; 1-2 FL OZ/A; REQUESTOR PREFERS A 6-DAY OR LESS PHI DUE TO FREQUENT FRUIT HARVESTING; NEED SUPPORT FOR A BLASTER OR KNAP SACK GROUND APPLIC; MFG WILL SUPPORT A 5-DAY PHI:08/16

**E/CS Data Requirements:** 3-5 GOOD TRIALS OVER 2 YEARS

**E/CS Research Comments:**

**IR-4 Residue Trial Plan:** 13-5; NEED 1 REG. 3/13 "RED A" TRIAL IN 2018 TO REPLACE A LOST TRIAL IN PR

**Comments:** NO SUPPORT FOR HANDGUN APPLICATIONS:06/08; MFG REQUIRES EFFICACY DATA BEFORE RESIDUE STUDY:06/09; KEY EXPORT MARKETS INCLUDE CANADA AND JAPAN:10/15; MFG RE-CONSIDERING SUPPORT:07/16; MFG SUPPORTS, BUT STILL NEED SOME EFFICACY DATA:08/16

**NER-EPA Region-FRD**

**NCR-EPA Region-FRD**

**SOR-EPA Region-FRD**

**WSR-EPA Region-FRD**

**CANADA-EPA Region-FRD**

18-FL4      Tannenbaum, Rebecca



## 2018 Tentative/Scheduled Studies Residue and Efficacy/Crop Safety (E/CS)

(Order by Crop Group, Commodity, Chemical)

Print Date: 2/22/2018

<u>PR #</u>	<u>LAB</u>	<u>PRIORITY</u>	<u>STUDY DIRECTOR</u>	<u>CHEMICAL (MFG)</u>	<u>COMMODITY</u>	<u>CROP GROUP</u>
P8274	-	A	HOMA	FLUAZINAM (ISK,SYNGEN)	PAPAYA	TROPICAL AND SUBTROPICAL, MEDIUM TO LARGE FRUIT, SMOOTH, INEDIBLE PEEL SUBGROUP (24B)

**Reason for Need:** ANTHRACNOSE

**Use Pattern: (PCR):** GROUND; 2 APPLIC; 14-DAY PHI; 1-2 FL OZ/A; REQUESTOR PREFERS A 6-DAY OR LESS PHI DUE TO FREQUENT FRUIT HARVESTING; NEED SUPPORT FOR A BLASTER OR KNAP SACK GROUND APPLIC; MFG WILL SUPPORT A 5-DAY PHI:08/16

**E/CS Data Requirements:** 3-5 GOOD TRIALS OVER 2 YEARS

**E/CS Research Comments:**

**IR-4 Residue Trial Plan:** 13-5; NEED 1 REG. 3/13 "RED A" TRIAL IN 2018 TO REPLACE A LOST TRIAL IN PR

**Comments:** NO SUPPORT FOR HANDGUN APPLICATIONS:06/08; MFG REQUIRES EFFICACY DATA BEFORE RESIDUE STUDY:06/09; KEY EXPORT MARKETS INCLUDE CANADA AND JAPAN:10/15; MFG RE-CONSIDERING SUPPORT:07/16; MFG SUPPORTS, BUT STILL NEED SOME EFFICACY DATA:08/16

**NER-EPA Region-FRD**

**NCR-EPA Region-FRD**

**SOR-EPA Region-FRD**

**WSR-EPA Region-FRD**

**CANADA-EPA Region-FRD**

18-FLP10 Crane, Dr. Jonathan H.



## 2018 Tentative/Scheduled Studies Residue and Efficacy/Crop Safety (E/CS)

(Order by Crop Group, Commodity, Chemical)

Print Date: 2/22/2018

<u>PR #</u>	<u>LAB</u>	<u>PRIORITY</u>	<u>STUDY DIRECTOR</u>	<u>CHEMICAL (MFG)</u>	<u>COMMODITY</u>	<u>CROP GROUP</u>
12283	18-MIR07	A	SAMOIL	FLONICAMID (FMC,ISK)	POMEGRANATE	TROPICAL AND SUBTROPICAL, MEDIUM TO LARGE FRUIT, SMOOTH, INEDIBLE PEEL SUBGROUP (24B)

**Reason for Need:** APHID; PLANT BUG (LYGUS), INCLUDING LEAF-FOOTED BUG; WHITEFLY

**Use Pattern: (PCR):** USE THE BELEAF 50SG PRODUCT; MAKE 1-3 FOLIAR APPLIC OF 0.089-0.133 LB AI/A BY GROUND OR AIR, 7-DAY INTERVAL, 14-DAY PHI; USE MINIMUM 50 GPA BY GROUND, 10 GPA BY AIR

**E/CS Data Requirements:** 3 TRIALS; HIGH RATE, LOW RATE, STANDARD, CONTROL

**E/CS Research Comments:**

**IR-4 Residue Trial Plan:** 10-4; 1 TRIAL IS A DECLINE

**Comments:** KEY EXPORT MARKETS INCLUDE KOREA, EU, JAPAN, CANADA:07/17; MFG SUPPORTS AND MAY ASSIST FINANCIALLY TO OFFSET STUDY COSTS:08/17; RESEARCHABLE, RESIDUE & E/CS DATA NEEDED

**NER-EPA Region-FRD**

**NCR-EPA Region-FRD**

**SOR-EPA Region-FRD**

**WSR-EPA Region-FRD**

**CANADA-EPA Region-FRD**

18-CA71	Kyser, Guy
(decline)	
18-CA72	Kyser, Guy
18-CA73	Leach, Nathan
18-CA74	Skiles, Keri





## 2018 Tentative/Scheduled Studies

### Residue and Efficacy/Crop Safety (E/CS)

*(Order by Crop Group, Commodity, Chemical)*

*Print Date: 2/22/2018*

<u>PR #</u>	<u>LAB</u>	<u>PRIORITY</u>	<u>STUDY DIRECTOR</u>	<u>CHEMICAL (MFG)</u>	<u>COMMODITY</u>	<u>CROP GROUP</u>
P12283	-	A	SAMOIL	FLONICAMID (FMC,ISK)	POMEGRANATE	TROPICAL AND SUBTROPICAL, MEDIUM TO LARGE FRUIT, SMOOTH, INEDIBLE PEEL SUBGROUP (24B)

**Reason for Need:** APHID; PLANT BUG (LYGUS), INCLUDING LEAF-FOOTED BUG; WHITEFLY

**Use Pattern: (PCR):** USE THE BELEAF 50SG PRODUCT; MAKE 1-3 FOLIAR APPLIC OF 0.089-0.133 LB AI/A BY GROUND OR AIR, 7-DAY INTERVAL, 14-DAY PHI; USE MINIMUM 50 GPA BY GROUND, 10 GPA BY AIR

**E/CS Data Requirements:** 3 TRIALS; HIGH RATE, LOW RATE, STANDARD, CONTROL

**E/CS Research Comments:**

**IR-4 Residue Trial Plan:** 10-4; 1 TRIAL IS A DECLINE

**Comments:** KEY EXPORT MARKETS INCLUDE KOREA, EU, JAPAN, CANADA:07/17; MFG SUPPORTS AND MAY ASSIST FINANCIALLY TO OFFSET STUDY COSTS:08/17; RESEARCHABLE, RESIDUE & E/CS DATA NEEDED

**NER-EPA Region-FRD**

**NCR-EPA Region-FRD**

**SOR-EPA Region-FRD**

**WSR-EPA Region-FRD**

**CANADA-EPA Region-FRD**

18-CAP13    Haviland, Dave  
18-CAP14    Haviland, Dave



# 2018 Tentative/Scheduled Studies

## Residue and Efficacy/Crop Safety (E/CS)

(Order by Crop Group, Commodity, Chemical)

Print Date: 2/22/2018

<u>PR #</u>	<u>LAB</u>	<u>PRIORITY</u>	<u>STUDY DIRECTOR</u>	<u>CHEMICAL (MFG)</u>	<u>COMMODITY</u>	<u>CROP GROUP</u>
P11754	-	A	HOMA	FLUXAPYROXAD + PYRACLOSTROBIN (BASF)	POMEGRANATE	TROPICAL AND SUBTROPICAL, MEDIUM TO LARGE FRUIT, SMOOTH, INEDIBLE PEEL SUBGROUP (24B)

**Reason for Need:** BLACK HEART, ALTERNARIA ROT, ROT OF FRUITS, COLLETOTRICHUM, CERCOSPORA, BOTRYOSPHERA, PILIDELIA, OTHER FUNGAL PATHOGENS OF FRUIT AND FOLIAGE

**Use Pattern: (PCR):** USE THE MERIVON PRODUCT; MAKE 2 FOLIAR APPLIC OF 6.2 FL OZ OF PROD/A, 14-21 DAY INTERVAL, 35-DAY PHI

**E/CS Data Requirements:** MFG REQUESTS THAT CERTAIN E/CS TESTING STANDARDS BE MET IN E/CS RESEARCH TO SUPPORT PRODUCT COMMERCIALIZATION; NEED SUFFICIENT

**E/CS Research Comments:** IN 2017 PROTOCOL: IN CA TRIALS, TESTING FOLIAR APPLIC OF MERVION AT LABEL RATE AND 2X LABEL RATE IN 20-100 GPA, WITH LATRON B-1956 ADJUVANT, ON THE SAME MAJOR VARIETIES THAT WERE TESTED IN 2016; MAKE 2 APPLIC, 14-DAY INTERVAL STARTING AT BLOOM; IN FL TRIALS, TESTING FOLIAR APPLIC OF MERVION AT LABEL RATE AND 2X LABEL RATE IN 20-100 GPA, WITH LATRON B-1956 ADJUVANT, ON THE SAME MAJOR VARIETIES THAT WERE TESTED IN 2016; MAKE 2 APPLIC, 14-DAY INTERVAL, WITH 2 TREATMENTS STARTING AT BLOOM AND THE OTHER 2 TREATMENTS STARTING SO THE LAST APPLIC IS 14 DAYS BEFORE HARVEST; EVALUATING CROP SAFETY AT LEAST 5 TIMES, AND FRUIT AND FOLIAR DISEASE CONTROL (INCIDENCE AND SEVERITY), ESPECIALLY FOR STYLAR AND BLACK HEART INFECTION AND ANTHRACNOSE INFECTION OF FRUIT, AS WELL AS FOLIAR SEVERITY OF ALTERNARIA SPP AND COLLETOTRICHUM SPP

**IR-4 Residue Trial Plan:** 3 10-4 (1 DECLINE); 1 PROCESSING (JUICE)

**Comments:** KEY EXPORT MARKETS INCLUDE KOREA, EU, RUSSIA; NO EFFECTIVE PRODUCT IS REGISTERED; THERE ARE NO TOLERANCES FOR EITHER AI:08/15; NEED THIS AS A ROTATIONAL PRODUCT IN A SUSTAINABLE DISEASE MANAGEMENT PROGRAM FOR FOLIAR AND FRUIT DISEASES IN SOUTHEAST STATES:09/15

NER-EPA Region-FRD

NCR-EPA Region-FRD

SOR-EPA Region-FRD

WSR-EPA Region-FRD

CANADA-EPA Region-FRD

18-CAP15 Michilaides, T.  
Fresno County  
18-CAP16 Michilaides, T.  
Kern County



## 2018 Tentative/Scheduled Studies

### Residue and Efficacy/Crop Safety (E/CS)

*(Order by Crop Group, Commodity, Chemical)*

*Print Date: 2/22/2018*

<b><u>PR #</u></b>	<b><u>LAB</u></b>	<b><u>PRIORITY</u></b>	<b><u>STUDY DIRECTOR</u></b>	<b><u>CHEMICAL (MFG)</u></b>	<b><u>COMMODITY</u></b>	<b><u>CROP GROUP</u></b>
07830	18-MIR01	A	BARNEY	ABAMECTIN (SYNGEN)	SUGAR APPLE	TROPICAL AND SUBTROPICAL, MEDIUM TO LARGE FRUIT, ROUGH OR HAIRY, INEDIBLE PEEL SUBGROUP (24C)

**Reason for Need:** MITES, LEAFMINER, THRIPS

**Use Pattern: (PCR):** 0.023 LB AI/A; 2 APPLIC AT A 30-DAY INTERVAL; 14-DAY PHI

**E/CS Data Requirements:**

**E/CS Research Comments:**

**IR-4 Residue Trial Plan:** 3/13-4; 1 TRIAL IS A DECLINE (2-YR STUDY); NEED 2 "RED A" TRIALS IN 2019

**Comments:** MFG WILL NOT SUPPORT:05/08; MFG CONFIRMED THIS USE IS NOW SUPPORTABLE:05/12; EPA (HOLD) CAUTION:08/14; EPA CAUTION:08/15; EPA CAUTION:08/16; EPA CAUTION: 08/17

**NER-EPA Region-FRD**

**NCR-EPA Region-FRD**

**SOR-EPA Region-FRD**

**WSR-EPA Region-FRD**

**CANADA-EPA Region-FRD**

18-FL107 Tannenbaum, Rebecca  
 18-FL108 Tannenbaum, Rebecca  
 (decline)



## 2018 Tentative/Scheduled Studies

### Residue and Efficacy/Crop Safety (E/CS)

*(Order by Crop Group, Commodity, Chemical)*

*Print Date: 2/22/2018*

<u>PR #</u>	<u>LAB</u>	<u>PRIORITY</u>	<u>STUDY DIRECTOR</u>	<u>CHEMICAL (MFG)</u>	<u>COMMODITY</u>	<u>CROP GROUP</u>
11292	17-FLR04	A	BARNEY	FLUMIOXAZIN (VALENT)	SUGAR APPLE	TROPICAL AND SUBTROPICAL, MEDIUM TO LARGE FRUIT, ROUGH OR HAIRY, INEDIBLE PEEL SUBGROUP (24C)

**Reason for Need:** PARTHENIUM WEED/RESIDUAL CONTROL

**Use Pattern: (PCR):** 2-6 OZ AI/A; 3-4 FOLIAR OR BANDED APPLIC DIRECTED TO WEEDS (MFG RECOMMENDS NO MORE THAN 24 OZ/A/YR); MIX WITH SYSTEMIC OR BURNDOWN HERBICIDE TO CONTROL MATURE PARTHENIUM; 30-90 DAY RE-TREATMENT INTERVAL (MFG RECOMMENDS 60 DAYS); 0-10 DAY PHI

**E/CS Data Requirements:** MFG REQUESTS 2X CROP SAFETY DATA, AND THAT THE LYCHEE E/CS PROTOCOL BE DUPLICATED, BUT MAKE SPRAY CROSS THE TRUNK SO THAT IN-RO'

**E/CS Research Comments:**

**IR-4 Residue Trial Plan:** 3/13-4, 1 DECLINE TRIAL (1 TRIAL IS COVERED BY TASC \$\$) (DOING 2 TRIALS IN 2017, 2 "RED A" TRIALS IN 2018)

**Comments:** EPA CAUTION:08/14; MFG ADDRESSING ISSUES:09/14; EPA CAUTION:08/16; RESIDUE & E/CS COMPLETE WITH ONGOING TRIALS:12/16;

**NER-EPA Region-FRD**

**NCR-EPA Region-FRD**

**SOR-EPA Region-FRD**

**WSR-EPA Region-FRD**

**CANADA-EPA Region-FRD**

18-FL114 Tannenbaum, Rebecca  
 18-FL115 Tannenbaum, Rebecca



## 2018 Tentative/Scheduled Studies

### Residue and Efficacy/Crop Safety (E/CS)

*(Order by Crop Group, Commodity, Chemical)*

*Print Date: 2/22/2018*

<u>PR #</u>	<u>LAB</u>	<u>PRIORITY</u>	<u>STUDY DIRECTOR</u>	<u>CHEMICAL (MFG)</u>	<u>COMMODITY</u>	<u>CROP GROUP</u>
P12110	-		DORSCHNER	INSECTICIDE (TBD)	PRICKLY PEAR CACTUS	TROPICAL AND SUBTROPICAL, CACTUS, INEDIBLE PEEL SUBGROUP (24D)

**Reason for Need:** FROM PR# 11964: COCHINEAL INSECTS (SUCKING INSECTS - HEMIPTERANS); WIND-BLOWN COCHINEAL INFESTATIONS (6, 2-MONTH GENERATIONS/YEAR) IN THE PAST 3 YEARS, LACK OF CONTROL FROM CARBARYL AND ACREAGE UNDER EXP USE PERMIT (FOR COCHINEAL CONTROL WITH SIVANTO) HAVE RESULTED IN 120 ACRES BEING BULLDOZED; WITH NEW CACTUS ORCHARDS TAKING 4 YEARS TILL FULL PRODUCTION, COCHINEALS HAVE RESULTED IN HUGE FINANCIAL LOSSES; WITH ONLY CARBARYL (2 APPLIC/YR) AND SIVANTO LABEL PENDING (2 APPLIC/YR), AT LEAST 4 MORE APPLIC/YR ARE NEEDED WITH INSECTICIDES FROM DIFFERENT CLASSES TO PROVIDE ECONOMIC CONTROL

**Use Pattern: (PCR):**

**E/CS Data Requirements:**

**E/CS Research Comments:** TESTING 2 RATES OF BOTH SULFOXAFLOL AND FLONICAMID, VS A STANDARD AND UNTREATED; MAKE FOLIAR APPLIC TIMED TO TREAT CRAWLER STAGE; MINIMUM 7-DAY RETREATMENT INTERVAL, IF NEEDED; COLLECTING DATA ON PEST CONTROL, CROP INJURY AND CROP RESPONSE

**IR-4 Residue Trial Plan:**

**Comments:** THIS PR# WAS CREATED AS A PPWS TO EVALUATE PERFORMANCE OF BOTH SULFOXAFLOL (SEE PR# 11964) AND FLONICAMID (SEE PR# 11966) UNDER A SINGLE PR#; SULFOXAFLOL WAS GIVEN AN "A" PRIORITY FOR RESIDUE WORK IN 2017, BUT THE DECISION WAS MADE THAT INSUFFICIENT EFFICACY DATA EXISTS TO SUPPORT THIS YET, SO A 2017 RESIDUE STUDY IS BEING DONE WITH FLONICAMID; IF SULFOXAFLOL PERFORMS EFFECTIVELY AND REGISTRATION IS DESIRED, PR 11964 WILL NEED TO GO THROUGH THE PRIORITIZATION PROCESS AGAIN:02/17

**NER-EPA Region-FRD**

**NCR-EPA Region-FRD**

**SOR-EPA Region-FRD**

**WSR-EPA Region-FRD**

**CANADA-EPA Region-FRD**

18-TBD      TBD-WSR  
(looking for Salinas Valley cooperators - tc)



# 2018 Tentative/Scheduled Studies Residue and Efficacy/Crop Safety (E/CS)

(Order by Crop Group, Commodity, Chemical)

Print Date: 2/22/2018

<b><u>PR #</u></b>	<b><u>LAB</u></b>	<b><u>PRIORITY</u></b>	<b><u>STUDY DIRECTOR</u></b>	<b><u>CHEMICAL (MFG)</u></b>	<b><u>COMMODITY</u></b>	<b><u>CROP GROUP</u></b>
11934	17-TIR01	A	JOLLY	AZOXYSTROBIN + CYPROCONAZOLE (SYNGEN)	COFFEE	MISC GROUP (99)

**Reason for Need:** COFFEE RUST (HEMILEIA VASTATRIX); NOT YET AN ISSUE IN HI, BUT NEED TO BE PREPARED

**Use Pattern: (PCR):** USE THE QUADRIS XTRA PRODUCT; MAKE 2-3 FOLIAR APPLIC OF 6.8 FL OZ PROD/A, 30-DAY INTERVAL, 30-DAY PHI; MAX 13.6 FL OZ/A/SEASON

**E/CS Data Requirements:**

**E/CS Research Comments:**

**IR-4 Residue Trial Plan:** 13-8 (3 TRIALS ARE COVERED BY TASC \$S) (DOING 5 TRIALS IN 2017, 3 "RED A" TRIALS IN 2018 [MAY NOT BE NEEDED - MFG HAS DATA FROM 5 TRIALS]); IN 2018 ONLY NEED 1 REG. 13 TRIAL, REPLACING A LOST PR TRIAL

**Comments:** TOLERANCES ARE AVAILABLE OUTSIDE THE US AND PUERTO RICO; IMPORT TOLERANCES ARE ESTABLISHED FOR BOTH AIS ON COFFEE IN THE US; MFG REQUESTS THAT THE USE PATTERN FOR THIS PROJECT SHOULD CONFORM TO WHAT IS ALREADY ESTABLISHED FOR USE ON COFFEE OUTSIDE THE US:06/16; CYPROCONAZOLE IS AN EPA CAUTION:08/16; MFG CONFIRMED THERE IS NO DECLINE OR PROCESSING DATA NEEDED; MFG HAS DATA FROM 5 TRIALS - IF THE USE PATTERN IN THIS STUDY IS THE SAME, THERE SHOULD BE NO NEED FOR ADDITIONAL "RED A" TRIALS IN 2018:11/16

**NER-EPA Region-FRD**

**NCR-EPA Region-FRD**

**SOR-EPA Region-FRD**

**WSR-EPA Region-FRD**

**CANADA-EPA Region-FRD**

18-PR311 Robles Vazquez, W.



## 2018 Tentative/Scheduled Studies

### Residue and Efficacy/Crop Safety (E/CS)

*(Order by Crop Group, Commodity, Chemical)*

Print Date: 2/22/2018

<b>PR #</b>	<b>LAB</b>	<b>PRIORITY</b>	<b>STUDY DIRECTOR</b>	<b>CHEMICAL (MFG)</b>	<b>COMMODITY</b>	<b>CROP GROUP</b>
11527	18-FLR03	A	SAMOIL	BIFENTHRIN (ADAMA,AMVAC,FMC)	COFFEE	MISC GROUP (99)

**Reason for Need:** COFFEE BERRY BORER, HYPOTHENEMUS HAMPEI; IS A RECENTLY INTRODUCED PEST WITH LIMITED CONTROL OPTIONS

**Use Pattern: (PCR):** 0.2-0.25 LB AI/A, MAX 0.5 LB AI/SEASON; 2 FOLIAR APPLIC/SEASON; RE-TREATMENT INTERVAL NOT LESS THAN 7 DAYS; 14-DAY PHI

**E/CS Data Requirements:** SUGGEST WE NEED GOOD RESULTS FROM A MINIMUM 3-4 FIELD TRIALS, HI AND PR; EFFICACY DATA REQUIREMENT IS NOW COMPLETE:09/16

**E/CS Research Comments:** NEED TO TARGET ADULTS EMERGING FROM BERRIES BEFORE THEY ENTER UNINFESTED BERRIES (ALL OTHER LIFE STAGES ARE PROTECTED INSIDE THE BERRIES); 2016 PERFORMANCE PROTOCOL: TEST 0.1 AND 0.2 LB AI/A RATES, VS A STANDARD, INCLUDING A NON-IONIC SURFACTANT; MAKE ONE APPLIC TO BERRIES WHICH ARE FREE OF BERRY BORERS; DATA ON CROP INJURY AND BORER CONTROL WILL BE COLLECTED

**IR-4 Residue Trial Plan:** 13-8; 1 TRIAL IS A DECLINE; 1 TRIAL IS FOR PROCESSING (GREEN BEAN, ROASTED, INSTANT); 2-YR STUDY; NEED 4 "RED A" TRIALS IN 2019 (2 IN PR, 2 IN HI) (3 TRIALS ARE COVERED BY TASC \$S, 2 IN 2018 AND 2 IN 2019)

**Comments:** EPA (HOLD) CAUTION:08/14; ADAMA & AMVAC WILL NOT SUPPORT; FMC OK:09/14; MFG NEEDS EFF DATA BEFORE SUPPORTING RESIDUE STUDY; MFG PREFERS HERO PRODUCT WITH ZETA-CYPERMETHRIN:07/15; EPA CAUTION:08/15; AT 2015 FUW, STAKEHOLDERS MADE THIS A "+" (H+) FOR THE 2016 PERFORMANCE PROGRAM:09/15; MFG CONFIRMED E/CS NEED IS SATISFIED, MAKING THIS RESIDUE-RESEARCHABLE:09/16; EPA CAUTION: 08/17

**NER-EPA Region-FRD**

**NCR-EPA Region-FRD**

**SOR-EPA Region-FRD**

**WSR-EPA Region-FRD**

**CANADA-EPA Region-FRD**

18-PR309 Robles Vazquez, W.  
18-PR310 Robles Vazquez, W.

18-HI150 Kam, James  
(processing)  
18-HI151 Kam, James  
(TASC \$)(decline)  
18-HI152 Kam, James  
(TASC \$)



## 2018 Tentative/Scheduled Studies

### Residue and Efficacy/Crop Safety (E/CS)

*(Order by Crop Group, Commodity, Chemical)*

*Print Date: 2/22/2018*

<b>PR #</b>	<b>LAB</b>	<b>PRIORITY</b>	<b>STUDY DIRECTOR</b>	<b>CHEMICAL (MFG)</b>	<b>COMMODITY</b>	<b>CROP GROUP</b>
P12390	-		DORSCHNER	CYANTRANILIPROLE (HGW86) (HSB)	CURRY LEAF	MISC GROUP (99)

**Reason for Need:** ASIAN CITRUS PSYLLID, THE VECTOR OF HUANGLONGBING

**Use Pattern: (PCR):** USE EXIREL OR VERIMARK PRODUCT; MAKE SOIL APPLIC OF 0.261-0.391 LB AI/A (MFG CONFIRMED THIS RATE RANGE IS FOR VERIMARK) VIA CHEMIGATION, 30-DAY INTERVAL, 1-DAY PHI; APPLY TO FIRST THIRD OF THE IRRIGATION CYCLE; DO NOT IRRIGATE FOR 24 HR AFTER APPLIC; PER MFG, BOTH EXIREL AND VERIMARK ARE EFFICACIOUS, BUT EXIREL (FOLIAR) HAS BEEN THE BEST OPTION, BASED ON USE RATES AND PRICE; FOR EXIREL, RESEARCHER INDICATES RATE RANGE IS 0.088-0.133 LB AI/A IN 100-150 GPA, WITH THE 0.133 LB AI RATE CONSIDERED BEST IN ORDER TO ENSURE EFFICACY:10/17

**E/CS Data Requirements:** 2 TRIALS; 1X, 2X, STANDARD, UNTREATED:10/17

**E/CS Research Comments:**

**IR-4 Residue Trial Plan:**

**Comments:** COMMODITY OF INTEREST IS INDIAN CURRY LEAVES; INSECTICIDES REGISTERED FOR CITRUS ARE FOR FRUIT CONSUMPTION, BUT WITH INDIAN CURRY LEAVES, MURRAYA KOENIGII, THE LEAVES ARE CONSUMED, NOT FRUIT; NO PRODUCTS ARE REGISTERED TO COVER LEAF CONSUMPTION:09/17; MFG (DUPONT HSB) SUPPORTS THIS USE, AND REQUIRES CROP SAFETY DATA, AS LEAVES ARE THE PARTS CONSUMED; THE AI IS EXCELLENT AGAINST CITRUS PSYLLID; RESIDUE PROJECT BEING CONDUCTED UNDER OTHER PR#: PR# 12402/MINT & PR# 12403/BASIL:10/17

**NER-EPA Region-FRD**

**NCR-EPA Region-FRD**

**SOR-EPA Region-FRD**

**WSR-EPA Region-FRD**

**CANADA-EPA Region-FRD**

18-CAP17    Mauk, Peggy A  
18-CAP18    Mauk, Peggy A





# 2018 Tentative/Scheduled Studies

## Residue and Efficacy/Crop Safety (E/CS)

*(Order by Crop Group, Commodity, Chemical)*

*Print Date: 2/22/2018*

<b><u>PR #</u></b>	<b><u>LAB</u></b>	<b><u>PRIORITY</u></b>	<b><u>STUDY DIRECTOR</u></b>	<b><u>CHEMICAL (MFG)</u></b>	<b><u>COMMODITY</u></b>	<b><u>CROP GROUP</u></b>
P12340	-	+	ARSENOVIC	HERBICIDE (TBD)	HEMP, INDUSTRIAL	MISC GROUP (99)

**Reason for Need:** WEEDS - THERE ARE NO REGISTERED HERBICIDES

**Use Pattern: (PCR):** NEED TO TEST VARIOUS PRODUCTS, FOLLOWING USE DIRECTIONS LABELED ON OTHER CROPS

**E/CS Data Requirements:** SUGGESTION IS FOR 4 TRIALS (NC, KY, WA); EXPECT 2 PROTOCOLS PER SITE: ONE IS A PRE APPLIC (PROBABLY SPIKE TIMING); SECOND IS AN EARLY PO

**E/CS Research Comments:**

**IR-4 Residue Trial Plan:**

**Comments:** CANADA PMC HAS DATA ON A NUMBER OF AIS:10/17

**NER-EPA Region-FRD**

**NCR-EPA Region-FRD**

**SOR-EPA Region-FRD**

**WSR-EPA Region-FRD**

**CANADA-EPA Region-FRD**

18-NCP02 Post, Dr. Angela R  
(preemergent)  
18-NCP03 Post, Dr. Angela R  
(postemergent)

18-WAP03 Schreiber, Mr. Alan (ADG  
(preemergent)  
18-WAP04 Schreiber, Mr. Alan (ADG  
(postemergent)



# 2018 Tentative/Scheduled Studies

## Residue and Efficacy/Crop Safety (E/CS)

(Order by Crop Group, Commodity, Chemical)

Print Date: 2/22/2018

<b>PR #</b>	<b>LAB</b>	<b>PRIORITY</b>	<b>STUDY DIRECTOR</b>	<b>CHEMICAL (MFG)</b>	<b>COMMODITY</b>	<b>CROP GROUP</b>
12346	18-CAR13	A	DORSCHNER	CYANTRANILIPROLE (HGW86) (HSB)	HOPS	MISC GROUP (99)

**Reason for Need:** HOP FLEA BEETLE, PSYLLIODES PUNCTULATUS - THERE ARE NO REGISTERED CONTROLS; WEEVILS - OTIORHYNCHUS SPP (BLACK VINE WEEVIL)

**Use Pattern: (PCR):** USE THE EXIREL OR VERIMARK PRODUCT; FOR FOLIAR USE - USING EXIREL FOR FLEA BEETLES, MAKE APPLIC OF 0.065-0.133 LB AI/A (10-20.5 FL OZ/A), 5-DAY INTERVAL, 1-DAY PHI, NO MORE THAN 0.4 LB AI/A/SEASON; FOR SOIL USE - USING VERIMARK VIA DRIP IRRIGATION (CHEMIGATION) FOR WEEVILS, MAKE APPLIC OF 0.088-0.130 LB AI/A (6.75-13.5 FL OZ/A), 2 APPLIC, 1-DAY PHI, NO MORE THAN 0.4 LB AI/A/SEASON

**E/CS Data Requirements:**

**E/CS Research Comments:**

**IR-4 Residue Trial Plan:** 5 11-3 12 (NAFTA SITES); 1 TRIAL IS A DECLINE

**Comments:** KEY EXPORT MARKETS INCLUDE EU, JAPAN, CANADA, CODEX; THE EU HOP INDUSTRY ALSO CONSIDERS THIS AI A HIGH PRIORITY FOR REGISTRATION, AND THE USE PATTERN PROVIDED HERE INCLUDES THE EU PROPOSED DIRECTIONS FOR USE:08/17; POTENTIAL JOINT PROJECT WITH CANADA:10/17; MFG APPROVED, RESIDUE AND PERFORMANCE DATA NEEDED, AT SEPT 2017 FUW, AND IN WRITING:01/18

**NER-EPA Region-FRD**

**NCR-EPA Region-FRD**

**SOR-EPA Region-FRD**

**WSR-EPA Region-FRD**

**CANADA-EPA Region-FRD**

18-WI384 Chapman, Scott

18-ID164 Meeks, Mr. Will  
 18-OR304 Sturman, Peter  
 18-WA367 Peng, Wilson  
 (decline)  
 18-WA368 Peng, Wilson

18-QC315 Cloutier, Dominic



# 2018 Tentative/Scheduled Studies

## Residue and Efficacy/Crop Safety (E/CS)

(Order by Crop Group, Commodity, Chemical)

Print Date: 2/22/2018

<b><u>PR #</u></b>	<b><u>LAB</u></b>	<b><u>PRIORITY</u></b>	<b><u>STUDY DIRECTOR</u></b>	<b><u>CHEMICAL (MFG)</u></b>	<b><u>COMMODITY</u></b>	<b><u>CROP GROUP</u></b>
P12346	-	A	DORSCHNER	CYANTRANILIPROLE (HGW86) (HSB)	HOPS	MISC GROUP (99)

**Reason for Need:** HOP FLEA BEETLE, PSYLLIODES PUNCTULATUS - THERE ARE NO REGISTERED CONTROLS; WEEVILS - OTIORHYNCHUS SPP (BLACK VINE WEEVIL)

**Use Pattern: (PCR):** USE THE EXIREL OR VERIMARK PRODUCT; FOR FOLIAR USE - USING EXIREL FOR FLEA BEETLES, MAKE APPLIC OF 0.065-0.133 LB AI/A (10-20.5 FL OZ/A), 5-DAY INTERVAL, 1-DAY PHI, NO MORE THAN 0.4 LB AI/A/SEASON; FOR SOIL USE - USING VERIMARK VIA DRIP IRRIGATION (CHEMIGATION) FOR WEEVILS, MAKE APPLIC OF 0.088-0.130 LB AI/A (6.75-13.5 FL OZ/A), 2 APPLIC, 1-DAY PHI, NO MORE THAN 0.4 LB AI/A/SEASON

**E/CS Data Requirements:**

**E/CS Research Comments:**

**IR-4 Residue Trial Plan:** 5 11-3 12 (NAFTA SITES); 1 TRIAL IS A DECLINE

**Comments:** KEY EXPORT MARKETS INCLUDE EU, JAPAN, CANADA, CODEX; THE EU HOP INDUSTRY ALSO CONSIDERS THIS AI A HIGH PRIORITY FOR REGISTRATION, AND THE USE PATTERN PROVIDED HERE INCLUDES THE EU PROPOSED DIRECTIONS FOR USE:08/17; POTENTIAL JOINT PROJECT WITH CANADA:10/17; MFG APPROVED, RESIDUE AND PERFORMANCE DATA NEEDED, AT SEPT 2017 FUW, AND IN WRITING:01/18

**NER-EPA Region-FRD**

**NCR-EPA Region-FRD**

**SOR-EPA Region-FRD**

**WSR-EPA Region-FRD**

**CANADA-EPA Region-FRD**

18-WAP05 Walsh, Dr. Doug



# 2018 Tentative/Scheduled Studies

## Residue and Efficacy/Crop Safety (E/CS)

(Order by Crop Group, Commodity, Chemical)

Print Date: 2/22/2018

<b>PR #</b>	<b>LAB</b>	<b>PRIORITY</b>	<b>STUDY DIRECTOR</b>	<b>CHEMICAL (MFG)</b>	<b>COMMODITY</b>	<b>CROP GROUP</b>
12334	18-CAR12	A	DORSCHNER	CYFLUMETOFEN (BASF)	HOPS	MISC GROUP (99)

**Reason for Need:** TWO SPOTTED SPIDER MITE, TETRANYCHUS URTICAE KOCH (ACARI: TETRANYCHIDAE); NEEDED FOR RESISTANCE MANAGEMENT DUE TO OTHER FAILING MITICIDES

**Use Pattern: (PCR):** FROM THE ORIGINAL PCR (PR# 10954): 0.179 LB AI/A; 2 FOLIAR APPLIC; 7-DAY RE-TREATMENT INTERVAL; 14-DAY PHI; NEW USE PATTERN COMMENTS: WILL TEST THE SAME RATE AS IN ORIGINAL PR# 10954 (MFG NOW KNOWS HOW TO MAKE THIS USE RATE WORK); NEED TO MATCH THE GERMAN USE RATE TO HARMONIZE MRLS; FINAL USE PATTERN IS NOT SET, BUT WILL BE CONFIRMED IN PROTOCOL DEVELOPMENT; ME-TOO REQUESTOR SUGGESTS 0.18 LB AI/A OF NEALTA, 2 FOLIAR AIRBLAST APPLIC 14 DAYS APART AND A 7-DAY PHI; THE USE PATTERN MUST MATCH THE USE PATTERN BEING DEVELOPED IN THE EU; MFG WILL NOT SUPPORT LABELING BACK TO BACK APPLIC OF NEALTA - AT LEAST ANOTHER MOA WILL BE REQUIRED BETWEEN NEALTA APPLIC:08/17

**E/CS Data Requirements:** MFG REQUIRES MINIMUM 6 EFFICACY TRIALS AND 3 CROP SAFETY TRIALS (ON POPULAR WA VARIETIES) SPREAD OVER 2 YEARS; MFG HAS DRAFTED EF

**E/CS Research Comments:**

**IR-4 Residue Trial Plan:** 5 11-3 12; 1 TRIAL IS A DECLINE (1 TRIAL IS COVERED BY TASC \$\$)

**Comments:** KEY EXPORT MARKETS INCLUDE THE EU, JAPAN, OTHERS; PRIOR RESIDUE STUDY (PR# 10954) WITH THIS AI ON HOPS WAS CANCELED DUE TO EFFICACY CONCERNS WITH THE LOW RATE OF CYFLUMETOFEN TESTED; EFFICACY CONCERNS HAVE NOW BEEN SUFFICIENTLY RESOLVED TO ALLOW THE MFG TO SUPPORT THIS USE AND RESIDUE WORK; ADDITIONAL CROP SAFETY DATA ON KEY VARIETIES IS REQUIRED, AND FURTHER EFFICACY NEEDS ARE BEING DETERMINED BY THE MFG:08/17; RESEARCHABLE, RESIDUE & E/CS DATA NEEDED

**NER-EPA Region-FRD**

**NCR-EPA Region-FRD**

**SOR-EPA Region-FRD**

**WSR-EPA Region-FRD**

**CANADA-EPA Region-FRD**

18-WI382      Chapman, Scott

18-ID161      Meeks, Mr. Will  
 18-OR302      Sturman, Peter  
 (decline)  
 18-WA362      Peng, Wilson  
 (TASC \$)  
 18-WA363      Peng, Wilson



# 2018 Tentative/Scheduled Studies

## Residue and Efficacy/Crop Safety (E/CS)

*(Order by Crop Group, Commodity, Chemical)*

Print Date: 2/22/2018

<b>PR #</b>	<b>LAB</b>	<b>PRIORITY</b>	<b>STUDY DIRECTOR</b>	<b>CHEMICAL (MFG)</b>	<b>COMMODITY</b>	<b>CROP GROUP</b>
P12334	-	A	DORSCHNER	CYFLUMETOFEN (BASF)	HOPS	MISC GROUP (99)

**Reason for Need:** TWO SPOTTED SPIDER MITE, TETRANYCHUS URTICAE KOCH (ACARI: TETRANYCHIDAE); NEEDED FOR RESISTANCE MANAGEMENT DUE TO OTHER FAILING MITICIDES

**Use Pattern: (PCR):** FROM THE ORIGINAL PCR (PR# 10954): 0.179 LB AI/A; 2 FOLIAR APPLIC; 7-DAY RE-TREATMENT INTERVAL; 14-DAY PHI; NEW USE PATTERN COMMENTS: WILL TEST THE SAME RATE AS IN ORIGINAL PR# 10954 (MFG NOW KNOWS HOW TO MAKE THIS USE RATE WORK); NEED TO MATCH THE GERMAN USE RATE TO HARMONIZE MRLS; FINAL USE PATTERN IS NOT SET, BUT WILL BE CONFIRMED IN PROTOCOL DEVELOPMENT; ME-TOO REQUESTOR SUGGESTS 0.18 LB AI/A OF NEALTA, 2 FOLIAR AIRBLAST APPLIC 14 DAYS APART AND A 7-DAY PHI; THE USE PATTERN MUST MATCH THE USE PATTERN BEING DEVELOPED IN THE EU; MFG WILL NOT SUPPORT LABELING BACK TO BACK APPLIC OF NEALTA - AT LEAST ANOTHER MOA WILL BE REQUIRED BETWEEN NEALTA APPLIC:08/17

**E/CS Data Requirements:** MFG REQUIRES MINIMUM 6 EFFICACY TRIALS AND 3 CROP SAFETY TRIALS (ON POPULAR WA VARIETIES) SPREAD OVER 2 YEARS; MFG HAS DRAFTED EF

**E/CS Research Comments:**

**IR-4 Residue Trial Plan:** 5 11-3 12; 1 TRIAL IS A DECLINE (1 TRIAL IS COVERED BY TASC \$\$)

**Comments:** KEY EXPORT MARKETS INCLUDE THE EU, JAPAN, OTHERS; PRIOR RESIDUE STUDY (PR# 10954) WITH THIS AI ON HOPS WAS CANCELED DUE TO EFFICACY CONCERNS WITH THE LOW RATE OF CYFLUMETOFEN TESTED; EFFICACY CONCERNS HAVE NOW BEEN SUFFICIENTLY RESOLVED TO ALLOW THE MFG TO SUPPORT THIS USE AND RESIDUE WORK; ADDITIONAL CROP SAFETY DATA ON KEY VARIETIES IS REQUIRED, AND FURTHER EFFICACY NEEDS ARE BEING DETERMINED BY THE MFG:08/17; RESEARCHABLE, RESIDUE & E/CS DATA NEEDED

**NER-EPA Region-FRD**

**NCR-EPA Region-FRD**

**SOR-EPA Region-FRD**

**WSR-EPA Region-FRD**

**CANADA-EPA Region-FRD**

18-WIP04 Heider, Daniel J.  
(crop safety)

18-ORP06 O'Neill, Greg (OR)  
(cv. Willamette & Nugget-Crop Safety)  
18-WAP06 Walsh, Dr. Doug  
(cv. Cascade & Tomahawk-Crop Safety)  
18-WAP07 Walsh, Dr. Doug  
(mite efficacy)



## 2018 Tentative/Scheduled Studies

### Residue and Efficacy/Crop Safety (E/CS)

*(Order by Crop Group, Commodity, Chemical)*

*Print Date: 2/22/2018*

<b><u>PR #</u></b>	<b><u>LAB</u></b>	<b><u>PRIORITY</u></b>	<b><u>STUDY DIRECTOR</u></b>	<b><u>CHEMICAL (MFG)</u></b>	<b><u>COMMODITY</u></b>	<b><u>CROP GROUP</u></b>
12342	18-TBD-CAN	A	CANADA	PYDIFLUMETOFEN (FTH 545) (SYNGEN)	HOPS	MISC GROUP (99)

**Reason for Need:** POWDERY MILDEW

**Use Pattern: (PCR):** MAKE FOLIAR APPLIC; OTHER USE PATTERN INFO NEEDED FROM US HOPS COMMISSION

**E/CS Data Requirements:**

**E/CS Research Comments:**

**IR-4 Residue Trial Plan:** NAFTA SITES: 5 11-3 12; IS A PMC CANADA 2018 STUDY (CANADAAS SD); 1 TRIAL IS A DECLINE

**Comments:** KEY EXPORT MARKETS PER OTHER HOPS USES; THERE IS INTEREST IN CANADA FOR THIS USE ("A" PRIORITY FOR 2018):08/17; MFG SUPPORTS, RESIDUE AND E/CS DATA NEEDED; HQ INCLUDING THIS AS A "RED A" TO ADD THREE U.S. TRIALS (REGION 11) TO THE 2018 PRIORITY "A" CANADIAN STUDY TO SUPPORT THIS AS A JOINT NAFTA STUDY:09/17

**NER-EPA Region-FRD**

**NCR-EPA Region-FRD**

**SOR-EPA Region-FRD**

**WSR-EPA Region-FRD**

**CANADA-EPA Region-FRD**

18-ID162	Meeks, Mr. Will	18-BC7	Clodius, Markus
18-OR303	Sturman, Peter	18-ON286	Wismer, R.J.
18-WA364	Peng, Wilson	(decline)	
18-WA365	Peng, Wilson		



# 2018 Tentative/Scheduled Studies

## Residue and Efficacy/Crop Safety (E/CS)

*(Order by Crop Group, Commodity, Chemical)*

*Print Date: 2/22/2018*

<b>PR #</b>	<b>LAB</b>	<b>PRIORITY</b>	<b>STUDY DIRECTOR</b>	<b>CHEMICAL (MFG)</b>	<b>COMMODITY</b>	<b>CROP GROUP</b>
P12389	-		DORSCHNER	CYANTRANILIPROLE (HGW86) (HSB)	KAFFIR LIME CITRUS	MISC GROUP (99)

**Reason for Need:** ASIAN CITRUS PSYLLID, THE VECTOR OF HUANGLONGBING

**Use Pattern: (PCR):** USE EXIREL OR VERIMARK PRODUCT; MAKE SOIL APPLIC OF 0.261-0.391 LB AI/A (MFG CONFIRMED THIS RATE RANGE IS FOR VERIMARK) VIA CHEMIGATION, 30-DAY INTERVAL, 1-DAY PHI; APPLY TO FIRST THIRD OF THE IRRIGATION CYCLE; DO NOT IRRIGATE FOR 24 HR AFTER APPLIC; PER MFG, BOTH EXIREL AND VERIMARK ARE EFFICACIOUS, BUT EXIREL (FOLIAR) HAS BEEN THE BEST OPTION, BASED ON USE RATES AND PRICE; FOR EXIREL, RESEARCHER INDICATES RATE RANGE IS 0.088-0.133 LB AI/A IN 100-150 GPA, WITH THE 0.133 LB AI RATE CONSIDERED BEST IN ORDER TO ENSURE EFFICACY:10/17

**E/CS Data Requirements:** 2 TRIALS; 1X, 2X, STANDARD, UNTREATED:10/17

**E/CS Research Comments:**

**IR-4 Residue Trial Plan:**

**Comments:** COMMODITY OF INTEREST IS KAFFIR LIME LEAVES; INSECTICIDES REGISTERED FOR CITRUS ARE FOR FRUIT CONSUMPTION, BUT WITH KAFFIR LIME, CITRUS HYSTERIX, THE LEAVES ARE CONSUMED, NOT THE FRUIT; NO PRODUCTS ARE REGISTERED TO COVER LEAF CONSUMPTION:09/17; MFG (DUPONT HSB) SUPPORTS THIS USE, AND REQUIRES CROP SAFETY DATA, AS LEAVES ARE THE PARTS CONSUMED; THE AI IS EXCELLENT AGAINST CITRUS PSYLLID:10/17; RESIDUE PROJECT BEING CONDUCTED UNDER PR# 12401, CYANTRANILIPROLE/DILL:10/17

**NER-EPA Region-FRD**

**NCR-EPA Region-FRD**

**SOR-EPA Region-FRD**

**WSR-EPA Region-FRD**

**CANADA-EPA Region-FRD**

18-CAP19    Mauk, Peggy A  
18-CAP20    Mauk, Peggy A



# 2018 Tentative/Scheduled Studies

## Residue and Efficacy/Crop Safety (E/CS)

*(Order by Crop Group, Commodity, Chemical)*

*Print Date: 2/22/2018*

<b><u>PR #</u></b>	<b><u>LAB</u></b>	<b><u>PRIORITY</u></b>	<b><u>STUDY DIRECTOR</u></b>	<b><u>CHEMICAL (MFG)</u></b>	<b><u>COMMODITY</u></b>	<b><u>CROP GROUP</u></b>
12402	18-MIR12	A	DORSCHNER	CYANTRANILIPROLE (HGW86) (HSB)	MINT	MISC GROUP (99)

**Reason for Need:** SEE REASON FOR THIS PR# IN THE COMMENTS, AND THE NEED ON INDIAN CURRY LEAVES FOR ASIAN CITRUS PSYLLID IN PR# 12390

**Use Pattern: (PCR):** FROM REQUEST 12390 FOR INDIAN CURRY LEAVES: USE EXIREL OR VERIMARK PRODUCT; MAKE SOIL APPLIC OF 0.261-0.391 LB AI/A (MFG CONFIRMED THIS RATE RANGE IS FOR VERIMARK) VIA CHEMIGATION, 30-DAY INTERVAL, 1-DAY PHI; APPLY TO FIRST THIRD OF THE IRRIGATION CYCLE; DO NOT IRRIGATE FOR 24 HR AFTER APPLIC; PER MFG, BOTH EXIREL AND VERIMARK ARE EFFICACIOUS, BUT EXIREL (FOLIAR) HAS BEEN THE BEST OPTION, BASED ON USE RATES AND PRICE; FOR EXIREL, RESEARCHER INDICATES RATE RANGE IS 0.088-0.133 LB AI/A IN 100-150 GPA, WITH THE 0.133 LB AI RATE CONSIDERED BEST IN ORDER TO ENSURE EFFICACY:10/17

**E/CS Data Requirements:**

**E/CS Research Comments:**

**IR-4 Residue Trial Plan:** 5-2 11-3 (1 DECLINE); FRESH AND DRY MINT FROM ALL TRIALS; OIL FROM 2 TRIALS

**Comments:** PROJECT ADDED FOR THIS FUTURE REP CROP FOR CROP GROUP 25 (HERBS), WHICH WILL PROVIDE A TOLERANCE THAT WILL COVER USE ON INDIAN CURRY LEAVES (SEE PR# 12390):10/17

**NER-EPA Region-FRD**

**NCR-EPA Region-FRD**

**SOR-EPA Region-FRD**

**WSR-EPA Region-FRD**

**CANADA-EPA Region-FRD**

18-WI389 Chapman, Scott  
(decline)  
18-WI390 Chapman, Scott  
(oil)

18-ID168 Meeks, Mr. Will  
18-OR305 Sturman, Peter  
(reg 11)  
18-WA371 Peng, Wilson  
(oil)





## 2018 Tentative/Scheduled Studies Residue and Efficacy/Crop Safety (E/CS)

(Order by Crop Group, Commodity, Chemical)

Print Date: 2/22/2018

<b>PR #</b>	<b>LAB</b>	<b>PRIORITY</b>	<b>STUDY DIRECTOR</b>	<b>CHEMICAL (MFG)</b>	<b>COMMODITY</b>	<b>CROP GROUP</b>
12347	18-CAR14	A	SAMOIL	LAMBDA-CYHALOTHRIN (SYNGEN)	MINT	MISC GROUP (99)

**Reason for Need:** CATERPILLARS, INCLUDING EARLY SEASON CUTWORMS AND LATE SEASON ARMY WORMS

**Use Pattern: (PCR):** USE THE WARRIOR II WITH ZEON TECHNOLOGY PRODUCT; MAKE 1 FOLIAR SPRAY OF 0.03 LB AI/A PER CUTTING, THROUGH CENTER PIVOT IRRIGATION SYSTEMS, 7-DAY PHI; APPLY IN MINIMUM OF 10 GPA; APPLY NO MORE THAN 0.06 LB AI/A/SEASON; DO NOT APPLY TO MINT THAT HAS PASSED 10% BLOOM (MFG NEEDS TO CONFIRM ASPECTS OF THE REQUESTED USE PATTERN:09/17)

**E/CS Data Requirements:**

**E/CS Research Comments:**

**IR-4 Residue Trial Plan:** 5-2 11-3; 2 TRIALS ARE PROCESSING (OIL); 1 TRIAL IS A DECLINE

**Comments:** MFG SUPPORTS, RESIDUE ONLY, WITH THE UNDERSTANDING THAT THIS USE COULD ONLY BE SUBMITTED IF THE 3X SAFETY FACTOR IS REMOVED:09/17; EPA CAUTION:11/17

**NER-EPA Region-FRD**

**NCR-EPA Region-FRD**

**SOR-EPA Region-FRD**

**WSR-EPA Region-FRD**

**CANADA-EPA Region-FRD**

18-WI385 Chapman, Scott  
18-WI386 Chapman, Scott  
(processing)

18-ID165 Meeks, Mr. Will  
(decline)  
18-ID166 Meeks, Mr. Will  
18-WA369 Peng, Wilson  
(processing)



## 2018 Tentative/Scheduled Studies Residue and Efficacy/Crop Safety (E/CS)

(Order by Crop Group, Commodity, Chemical)

Print Date: 2/22/2018

<u>PR #</u>	<u>LAB</u>	<u>PRIORITY</u>	<u>STUDY DIRECTOR</u>	<u>CHEMICAL (MFG)</u>	<u>COMMODITY</u>	<u>CROP GROUP</u>
12247	18-YAR03	A	BARNEY	S-METOLACHLOR/METOLAC HLOR (SYNGEN,UPI)	QUINOA	MISC GROUP (99)

**Reason for Need:** BROAD SPECTRUM CONTROL OF GRASSES AND BROAD-LEAF WEEDS - THERE ARE LIMITED OPTIONS FOR QUINOA GROWERS

**Use Pattern: (PCR):** USE THE DUAL II MAGNUM PRODUCT; MAKE 1 PRE-EMERGENT (SOIL-APPLIED OR POST-EMERGENT [WHEN QUINOA IS AT 4 LEAF STAGE])  
APPLIC OF 0.63 LB AI/A, 70-90 DAY PHI

**E/CS Data Requirements:** MFG REQUESTS SUFFICIENT CROP SAFETY DATA TO SATISFY THE MFG AND CA-DPR:08/17; MFG REQUIRES 4-5 TRIALS, 1X AND 2X RATES (AND MIXTURE

**E/CS Research Comments:**

**IR-4 Residue Trial Plan:** 11/12-4 (GRAIN ONLY)

**Comments:** A CODEX MRL IS IMPORTANT; IR-4 TO EXAMINE IF SEVERAL TOLERANCES ON OTHER CROPS CAN BE USED IN A CHEMSAC PROPOSAL TO PROPOSE A QUINOA TOLERANCE WITHOUT THE NEED FOR RESIDUE WORK:05/17; POTENTIAL JOINT PROJECT WITH CANADA:10/17; DECISION WAS MADE THAT RESIDUE DATA ARE NEEDED TO ESTABLISH A QUINOA TOLERANCE, AND TO SUPPORT INT'L MRLS:01/18

**NER-EPA Region-FRD**

**NCR-EPA Region-FRD**

**SOR-EPA Region-FRD**

**WSR-EPA Region-FRD**

**CANADA-EPA Region-FRD**

18-CA58	Kyser, Guy
18-ID157	Meeks, Mr. Will
18-OR297	Sturman, Peter
18-OR298	Sturman, Peter
(5X)	
18-WA353	Peng, Wilson
18-WA*354	Harvey, John
18-WA*355	Harvey, John
(5X)	



# 2018 Tentative/Scheduled Studies Residue and Efficacy/Crop Safety (E/CS)

(Order by Crop Group, Commodity, Chemical)

Print Date: 2/22/2018

---

<u>PR #</u>	<u>LAB</u>	<u>PRIORITY</u>	<u>STUDY DIRECTOR</u>	<u>CHEMICAL (MFG)</u>	<u>COMMODITY</u>	<u>CROP GROUP</u>
P12247	-	A	BARNEY	S-METOLACHLOR/METOLAC HLOR (SYNGEN,UPI)	QUINOA	MISC GROUP (99)

**Reason for Need:** BROAD SPECTRUM CONTROL OF GRASSES AND BROAD-LEAF WEEDS - THERE ARE LIMITED OPTIONS FOR QUINOA GROWERS

**Use Pattern: (PCR):** USE THE DUAL II MAGNUM PRODUCT; MAKE 1 PRE-EMERGENT (SOIL-APPLIED OR POST-EMERGENT [WHEN QUINOA IS AT 4 LEAF STAGE])  
APPLIC OF 0.63 LB AI/A, 70-90 DAY PHI

**E/CS Data Requirements:** MFG REQUESTS SUFFICIENT CROP SAFETY DATA TO SATISFY THE MFG AND CA-DPR:08/17; MFG REQUIRES 4-5 TRIALS, 1X AND 2X RATES (AND MIXTURE

**E/CS Research Comments:**

**IR-4 Residue Trial Plan:** 11/12-4 (GRAIN ONLY)

**Comments:** A CODEX MRL IS IMPORTANT; IR-4 TO EXAMINE IF SEVERAL TOLERANCES ON OTHER CROPS CAN BE USED IN A CHEMSAC PROPOSAL TO PROPOSE A QUINOA TOLERANCE WITHOUT THE NEED FOR RESIDUE WORK:05/17; POTENTIAL JOINT PROJECT WITH CANADA:10/17; DECISION WAS MADE THAT RESIDUE DATA ARE NEEDED TO ESTABLISH A QUINOA TOLERANCE, AND TO SUPPORT INT'L MRLS:01/18

**NER-EPA Region-FRD**

**NCR-EPA Region-FRD**

**SOR-EPA Region-FRD**

**WSR-EPA Region-FRD**

**CANADA-EPA Region-FRD**

18-ORP07 Peachey, Ed  
18-CAP21 Hanson, Brad  
18-WA\*P08 Harvey, John  
18-WAP09 Miller, Mr. Tim (MT Vern)



# 2018 Tentative/Scheduled Studies

## Residue and Efficacy/Crop Safety (E/CS)

(Order by Crop Group, Commodity, Chemical)

Print Date: 2/22/2018

---

<u>PR #</u>	<u>LAB</u>	<u>PRIORITY</u>	<u>STUDY DIRECTOR</u>	<u>CHEMICAL (MFG)</u>	<u>COMMODITY</u>	<u>CROP GROUP</u>
09871	18-CAR05	A	ARSENOVIC	ETHALFLURALIN (GOWAN)	STEVIA	MISC GROUP (99)

**Reason for Need:** WEEDS; FROM PR 09873: NEED A PREPLANT INCORPORATED USE PATTERN

**Use Pattern: (PCR):** 1.5-4.0 PT/A; GROUND APPLIC

**E/CS Data Requirements:**

**E/CS Research Comments:**

**IR-4 Residue Trial Plan:** ANY 4 TRIALS (DRIED LEAF)

**Comments:** MFG WILL NOT SUPPORT:07/09; MFG (GOWAN) PREFERS THIS AI INSTEAD OF TRIFLURALIN/STEVIA PR# 09873 WHICH WAS THE "A" PRIORITY AT THE 2017 FUW:10/17

**NER-EPA Region-FRD**

18-NJ235 Fisher, Jennifer

**NCR-EPA Region-FRD**

**SOR-EPA Region-FRD**

18-NC212 Batts, Roger B.  
18-NC337 Batts, Roger B.

**WSR-EPA Region-FRD**

18-CA21 Ennes, D. (Kearney)  
18-CA22 Watkins, S.

**CANADA-EPA Region-FRD**



## 2018 Tentative/Scheduled Studies Residue and Efficacy/Crop Safety (E/CS) (Order by Crop Group, Commodity, Chemical)

Print Date: 2/22/2018

---

	NER	NCR	SOR	WSR	CANADA
ARS Total:		11	21	25	
Region Total:	44	76	95	189	21
Total:	44	87	116	214	21

**Grand Trial Total: 482**

**Total # of PRs: 90**  
**Total # Chemical: 53**  
**Total # Commodity: 66**